


RIUNIONE SCIENTIFICA ANNUALE DEL GRUPPO DI ALGOLOGIA DELLA SOCIETA' BOTANICA ITALIANA (SBI)

Amendolara, 11-12 novembre 2022



Stazione Zoologica Anton Dohrn - Sede Calabria



deltek



Società Botanica Italiana
Gruppo di Algologia
Riunione Scientifica Annuale
Amendolara, 11-12 novembre 2022
Stazione Zoologica Anton Dohrn – Sede Calabria

Comitato organizzatore:
Chiara Pennesi e Chiara Giommi

Staff di supporto: Mario De Stefano, Claudia Ciniglia, Concetta Auciello, Luciana Barra, Valentina Costa, Paolo D'Ambrosio, Roberto Firmamento, Christian Galasso, Martina Genovese, Danilo Malara, Chiara Molino, Mirko Mutalipassi, Daniela Pica, Nadia Ruocco, Claudia Scianna, Serena Scozzafava, Teresa Manuela Vicchio.

Si ringraziano:
il Comune di Amendolara per aver patrocinato l'evento,
Deltek per aver finanziato il kit congressuale.



Programma

Venerdì, 11 novembre 2022

- 8.30 Inizio registrazione dei partecipanti
- 9.15 Saluti di benvenuto e introduzione ai lavori
- 10.00 Relazione ad invito:
Prof. Christopher S. Lobban, University of Guam
What if we assume everything is NOT everywhere?

Sessione Microalghe I - Moderatore: Rossella Pistocchi

- 10.30 **M. De Stefano, C. Auciello, M. Iovinella, C. Ciniglia, C. Pennesi**
From Antarctica to Tropics, a long term study on biodiversity and ecological role of benthic diatom communities
- 10.45 **R. Trentin, E. Moschin, A. Duarte Lopes, S. Schiaparelli, L. Custódio, I. Moro**
Molecular, morphological and chemical diversity of two new species of Antarctic diatoms, *Craspedostauros ineffabilis* sp. nov. and *Craspedostauros zucchellii* sp. nov.
- 11.00 **R. Bentivogli, L. Pezzolesi, N. Caputo, F. Costantini, J. Fondo, R. Pistocchi**
Morphological vs. innovative DNA metabarcoding approach for the phytoplankton monitoring in an Adriatic coastal lagoon (cascade project)

11.15 - 11.45 Coffee break

Sessione Macroalghe - Moderatore: Fabio Rindi

- 11.45 **G. Denti, A. Petrocelli, E. Cecere, G. Fanelli, L. Papa, F. Rubino**
Further evidence of the presence of *Caulerpa cylindracea* (Bryopsidales, Chlorophyta) and *Asparagopsis* sp. (Bonnemaisoniales, Rhodophyta) in the Taranto seas (Northern Ionian Sea, Mediterranean Sea), two non-indigenous species (NIS) of macroalgae
- 12.00 **D. Lenzo, A. Bartual, B. Úbeda, E. Zubía, B. Cuevas, M.A. Colangelo, A. Pasteris, L. Pezzolesi**
Allelopathic effects in the invasive macroalga *Rugulopterix okamurae* (Phaeophyceae):



potential role of the newly discovered compound Dilkamural

- 12.15 **D. Oliva, A. Piro, M. Carbone, E. Mollo, F. Scarcelli, V. Osso, D.M. Nisticò, S. Mazzuca**
Study of the effects of metabolites of invasive *Caulerpa* spp. on *Posidonia oceanica* seedling growth: unexpected effects of the vehicle (DMSO)
- 12.30 **A. Tursi, F. Mastrototaro, G. Chimienti, A. Bottalico**
Algal assemblages' composition of two semi-submerged marine caves at Tremiti Islands Marine Protected Area (Italy, Adriatic Sea)
- 12.45 **S. Kaleb, E. Descourvières, V. Bandelj, F. Gianni, M. Orlando-Bonaca, A. Falace**
New insights into the reproductive strategies of the endangered *Fucus virsoides* useful for implementing restoration measures

13.00– 15.00 Pausa pranzo

Sessione Microalghe II - Moderatore: Stefano Accoroni

- 15.00 **M. Marieschi, G. Molinari, S. Savi, R. Bolpagni, A. Torelli**
Search of freshwater algal species with adaptive mechanisms to Cr (VI) stress
- 15.15 **G. Molinari, M. Marieschi, R. Bolpagni, A. Torelli**
Interaction between OAS-TL and desulphydrase activity in Cr (VI) tolerance in the freshwater green alga *Scenedesmus acutus*
- 15.30 **L. Mollo, F. Drigo, A. Norici**
Effetti di composti fenolici presenti in acque di vegetazione sulle microalghe
- 15.45 **C. Auciello, C. Pennesi, C. Ciniglia, M. Iovinella, L. Rabaoui, M. De Stefano**
Benthic diatom communities on seagrass and corals in Arabian Gulf: floristic analysis, population dynamics and estimate of contribute to the primary production of the basin
- 16.00 **M. Blasio, G.V. de Klerk, P. Marotta, A. Cutignano, G. Trancone, G. Dupont, A. Sardo, P. De Luca, S. Papirio, S Balzano**
Potential of *Nannochloropsis oceanica* for bioremediation
- 16.15 **S. D'Arpa, M.R. Vadrucci, P.C. Ciciriello, E. Stanca, A. Pastorelli, N. Ungaro**
A photo guide of harmful algae along coastal areas of Apulian Region



16:30 - 17.00 Coffee break

Sessione Microalghe III - Moderatore: Marina Montresor

17.00 **V. Natali, T. Cibic**

Valutazione dell'effetto inibitorio di materiali legnosi modificati o trattati con impregnanti di nuova generazione sullo sviluppo della comunità microfitobentonica nella laguna di Grado (nord Adriatico)

17.15 **F. Neri, M. Ubaldi, T. Romagnoli, S. Accoroni, C. Totti**

Phytoplankton indicators for GES assessment: a case study in the N Adriatic Sea

17.30 **A. Pelusi, L. Ambrosino, M. Miraldo, M.L. Chiusano, A. Rogato, M.I. Ferrante, M. Montresor**

Espressione genica durante la formazione di spore nella diatomea marina *Chaetoceros socialis*

17.45 **L. Roselli, S. Bevilacqua, A. Terlizzi**

Trait-based approach for rapid and effective characterization of phytoplankton patterns

18.00 **G. Furnari**

Amenità fisiologiche e non solo

18.15 **ASSEMBLEA DEI SOCI**

20.30 Cena Sociale

Sabato, 12 novembre 2022

Sessione Microalghe IV - Moderatore: Laura Pezzolesi

9.30 **K. Sciuto, A.A. Sfriso, M.A. Wolf, A. Sfriso, E. Moschin, C. Munari, M. Mistri, I. Moro**

Dati preliminari su una microalga verde isolata dal Lido di Spina (Comacchio, Ferrara)

9.45 **M. Ubaldi, S. Bacchiocchi, M. Siracusa, F. Neri, T. Romagnoli, C. Totti, S. Accoroni**

Effect of salinity on growth and microcystin production by the cyanobacterium



Planktothrix rubescens

- 10.00 **L. Barra, S. Balzano, M. Moros, A. Smerilli, E. Cavalletti, I. Percopo, P. Chiaiese, A. Sardo**
Potential of a green algal strain collected from the Sarno River mouth for heavy metal bioremediation
- 10.15 **M. Simonazzi, L. Pezzolesi, T.K. Shishido, D.P. Fewer, R. Pistocchi**
Investigation of potentially toxic metabolites in a newly isolated freshwater cyanobacterium of the family Pseudanabaenaceae through a genomic approach
- 10.30 **M. Iovinella, C. Ciniglia, S.J. Davis**
Organelle phylogenomics unveil the diverging evolution of 6 lineages in *Galdieria sulphuraria* (Cyanidiophyceae)

10.45 - 11.15 Coffee break

Sessione: Microalge V - Moderatore Chiara Pennesi

- 11.15 **M. Mutualipassi, V. Mazzella, N. Ruocco, V. Zupo**
Halomicronema metazoicum, la storia di un cianobatterio, del suo ambiente e dei suoi vicini
- 11.30 **I. Orefice, V. Di Dato, G. Romano**
Induzione di stadi di resistenza nella diatomea marina *Thalassiosira rotula*
- 11.45 **G. Zoffoli, C. Samorì, M. Simonazzi, R. Pistocchi, L. Pezzolesi**
Production of the biopolymer polyhydroxybutyrate (PHB) from the Chlorophyceae *Desmodesmus communis*

12.00 Arrivederci!



Sessione Microalghe I

Moderatore: Rossella Pistocchi



From Antarctica to Tropics, a long term study on biodiversity and ecological role of benthic diatom communities

M. De Stefano, C. Auciello, M. Iovinella, C. Ciniglia, C. Pennesi

In Polar and tropical marine waters, microalgal communities associated to macrophytes, seagrasses, and invertebrates can be often more productive than phytoplanktonic ones (Colijn & De Jonge, 1984; Sundbäck & Jönsson, 1988). These benthic microalgal assemblages, known as "microphytobenthos" are mainly represented by diatoms (Bacillariophyta) (Mac Intyre et al., 1996). and specifically by a well-circumscribed group of "pennate" diatoms genera that can be considered adapted to this peculiar epibenthic life habitus. However, despite to their crucial ecological role in the trophic chain, the biodiversity of epibenthic diatom communities worldwide is poorly studied and greatly underestimated. Floristic studies at species or intraspecific level are rare since most of these species exhibit mean dimensions lower than 20 micrometers with taxonomic features often below the solving power of the light microscope. Therefore, a floristic and ecological analysis of these communities needs scanning electron microscopy (SEM) as common approach. In the framework of an international project aimed to decrypt and document the hidden biodiversity of epibenthic diatom assemblages worldwide, we will report the results of comparative analysis, in term of specie richness, community structure, and ecological role, within Antarctic and Tropical diatom communities associated to macrophytes and different invertebrates including gastropods, crustaceans bryozoans and bivalves. Our results demonstrate that either in Polar or Tropical regions, significant benthic diatom communities can be hosted by macrophytes whereas those on invertebrates are generally poor in term of species and relative abundances. Specifically, cell densities of seagrasses diatom communities resulted more than one hundred times higher than those of invertebrates with mean values approximately of 6000-7000 cell/mm² (up to 13000 cell/mm²) in the former vs 100-300 cell/mm² in the latter. Noteworthy is that the specie composition of seagrasses diatom communities in both biogeographic regions was characterized by the almost full dominance of adnate genus *Cocconeis* (represented by four species) exhibiting cell densities mean values more than one hundred times higher than those of all the other diatom genera present in the communities. *Cocconeis* predominance in epibenthic diatom assemblages was been also confirmed by previous similar studies on Mediterranean and indo-pacific regions thus confirming the crucial ecological role of such genera in all epibenthic diatom communities worldwide.

Letteratura citata

- Colijn, F., De Jonge, V.N. (1984). Primary production of microphytobenthos in the Ems-Dollard estuary. Mar. Ecol. Prog. Ser. 14: 185-196.
 MacIntyre, H.L., Geider, R.J., Miller, D.C. (1996). Microphytobenthos: the ecological role of the secret garden of unvegetated, shallowwater marine habitats. I. Distribution, abundance and primary production. *Estuaries*. 19: 186–201.
 Sundbäck, K., Jönsson, B. (1988). Microphytobenthic productivity and biomass in sublittoral sediments of a stratified bay, south-eastern Kattegat. *J. Exp. Mar. Biol. Ecol.* 122: 63-81.

AUTORI

Mario De Stefano (mario.destefano@unicampania.it); Autore di riferimento), Concetta Auciello (concetta.auciello@unicampania.it), Manuela Iovinella (manuela.iovinella@unicampania.it), Claudia Ciniglia (claudia.ciniglia@unicampania.it), Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "L. Vanvitelli", via Vivaldi 43, 81100 Caserta, Italia.
 Chiara Pennesi (chiara.penesi@szn.it), Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn, C.da Torre Spaccata, Località Torre Spaccata 87071 Amendolara (CS), Italia.



Molecular, Morphological and Chemical Diversity of two new species of Antarctic Diatoms, *Craspedostauros ineffabilis* sp. nov. and *Craspedostauros zucchellii* sp. nov.

R. Trentin, E. Moschin, A. Duarte Lopes, S. Schiaparelli, L. Custódio, I. Moro

The current study focuses on the biological diversity of two strains of Antarctic diatoms (strains IMA082A and IMA088A) collected and isolated from the Ross Sea (Antarctica) during the XXXIV Italian Antarctic Expedition. Both species presented the typical morphological characters of the genus *Craspedostauros*: ciliate areolae, two 'fore-and-aft' chloroplasts and a narrow 'stauros'. This classification is congruent with the molecular phylogeny based on the concatenated 18S rDNA-*rbcL-psbC* alignment, which showed that these algae formed a monophyletic lineage including six taxonomically accepted species of *Craspedostauros*. Since the study of the evolution of this genus and of others raphe-bearing diatoms with a 'stauros' is particularly challenging and their phylogeny is still debated, we tested alternative tree topologies to evaluate the relationships among these taxa. Metabolic fingerprinting approach was implemented for the assessment of the chemical diversity of IMA082A and IMA088A. In conclusion, combining: (1) traditional morphological features used in diatoms identification, (2) phylogenetic analyses of the small subunit rDNA (18S rDNA), *rbcL* and *psbC* genes, and (3) metabolic fingerprint, we described the strains IMA082A and IMA088A as *Craspedostauros ineffabilis* sp. nov. and *Craspedostauros zucchellii* sp. nov. as new species, respectively.

AUTORI

Riccardo Trentin, (riccardo.trentin.2@studenti.unipd.it; Autore di riferimento) Department of Biology, University of Padova, 35131 Padova, Italy. Emanuela Moschin, Department of Biology, University of Padova, 35131 Padova, Italia.
 André Duarte Lopes, Department of Chemistry and Pharmacy, FCT, University of the Algarve, 8005-039 Faro, Portugal; Department of Chemistry and Pharmacy, FCT, University of the Algarve, 8005-039 Faro, Portogallo.
 Stefano Schiaparelli, Department of Earth, Environment and Life Sciences, University of Genoa, 16132 Genova, Italy; Italian National Antarctic Museum (MNA, Section of Genoa), University of Genoa, 16132 Genova, Italia.
 Luísa Custódio, Centre of Marine Sciences, Faculty of Sciences and Technology, University of Algarve, 8005-139 Faro, Portugal
 Isabella Moro (isabella.moro@unipd.it), Department of Biology, University of Padova, 35131 Padova, Italia.



Morphological vs. innovative DNA metabarcoding approach for the phytoplankton monitoring in an Adriatic coastal lagoon (cascade project)

R. Bentivogli, L. Pezzolesi, N. Caputo, F. Costantini, J. Fondo, R. Pistocchi

The INTERREG project CASCADE (CoASTal and marine water integrated monitoring systems for ecosystems proteCtion AnD managemEnt) aims to assess and to evaluate the quality and the vulnerability of coastal ecosystems, with the final purpose to enhance the knowledge, restore endangered species and support an integrated management of coastal areas. Microalgae have a crucial role in water management since they represent the mainstay of coastal ecosystems. Given that algal blooms have become a widespread phenomenon across the world (Wurtsbaugh et al., 2019) and established that traditional monitoring methods based on the experience of the surveyor are not appropriate for a large-scale measurement of phytoplankton, new methodologies are needed to overcome the issue and face the diversified factors driving algal blooms (Wood et al., 2013). As part of the CASCADE project, during the last year a monthly monitoring campaign has been carried out in a coastal lagoon situated in the northern Adriatic Sea (Sacca di Goro, FE) to assess the planktonic communities and evaluate the ecological status of the area. The analysis of the phytoplankton communities relied on qualitative and quantitative evaluation of the species present using a morphological analysis (microscope), however, an innovative technique, mainly the environmental DNA metabarcoding using Oxford Nanopore MinIon sequencing, has been tested during seasonal samplings and compared with the microscope analysis to better define phytoplankton dynamics in a coastal lagoon characterized by peculiar hydrological trends. This innovating technique has enabled to detect new multiple, cryptic and picoplanktonic taxa that were not identified using microscope and, in addition, it was found to be effective in the identification of harmful algal bloom species and non-indigenous species in this shallow water lagoon. DNA metabarcoding has been largely applied in the last decade as an instrument to fulfil ecosystem biotic information (Brown et al., 2022). Although it represents a powerful tool for valuable insights, it still needs to be improved, specifically in the implementation of the barcoding libraries, thus it cannot yet be treated as a complete substitute for the traditional monitoring methods.

Letteratura citata

- Brown, P. D., Craine, J. M., Richards, D., Chapman, A., & Marden, B. (2022). DNA metabarcoding of the phytoplankton of Great Salt Lake's Gilbert Bay: Spatiotemporal assemblage changes and comparisons to microscopy. *Journal of Great Lakes Research*, 48(1), 110-124.
- Wood, S. A., Smith, K. F., Banks, J. C., Tremblay, L. A., Rhodes, L., Mountfort, D., ... & Pochon, X. (2013). Molecular genetic tools for environmental monitoring of New Zealand's aquatic habitats, past, present and the future. *New Zealand Journal of Marine and Freshwater Research*, 47(1), 90-119.
- Wurtsbaugh, W. A., Paerl, H. W., & Dodds, W. K. (2019). Nutrients, eutrophication and harmful algal blooms along the freshwater to marine continuum. *Wiley Interdisciplinary Reviews: Water*, 6(5), e1373.

AUTORI

Riccardo Bentivogli, (riccardo.bentivogli4@unibo.it; Autore di riferimento), Interdepartmental Centre for Research in Environmental Sciences – CIRSA, University of Bologna, Via Sant'Alberto 163, 48123, Ravenna, Italia.
 Laura Pezzolesi, Federica Costantini, Rossella Pistocchi, Interdepartmental Centre for Research in Environmental Sciences – CIRSA, University of Bologna, Via Sant'Alberto 163, 48123, Ravenna, Italy; Department of Biological, Geological and Environmental Sciences – BIGeA, University of Bologna, Ravenna Campus, Via Sant'Alberto 163, 48123, Ravenna, Italia.
 Nicolè Caputo, Interdepartmental Centre for Research in Environmental Sciences – CIRSA, University of Bologna, Via Sant'Alberto 163, 48123, Ravenna, Italy; Marine and Freshwater Research Centre (MFRC), Atlantic Technological University (ATU), Dublin Road, Galway City, Republic of Ireland (present address).



RIUNIONE SCIENTIFICA ANNUALE
Amendolara, 11-12 novembre 2022



Joyce Rukia Fondo, Department of Biological, Geological and Environmental Sciences – BIGeA, University of Bologna,
Ravenna Campus, Via Sant'Alberto 163, 48123, Ravenna, Italia.





Sessione Macroalge

Moderatore: Fabio Rindi



Further evidence of the presence of *Caulerpa cylindracea* (Bryopsidales, Chlorophyta) and *Asparagopsis* sp. (Bonnemaisoniales, Rhodophyta) in the Taranto seas (Northern Ionian Sea, Mediterranean Sea), two non-indigenous species (NIS) of macroalgae

G. Denti, A. Petrocelli, E. Cecere, G. Fanelli, L. Papa, F. Rubino

The introduction of non-indigenous species (NIS), organisms arriving from beyond their natural (past or present) geographical region and outside of their natural dispersal potential, represents one of the most important threats for biodiversity. This arrival in many cases leads to detrimental community-level modifications and, consequently, to alterations of ecosystem functioning together with socio-economic dimension that is unavoidably affected (Tsirintanis et al., 2022). 1.288 trillion \$ are reported as the minimum value of the global economic loss related to biological invasion assessed in the past few decades (1970-2017). The Mediterranean Sea is considered a hotspot for biological invasions counting more than 1000 registered non-indigenous species; of these, seaweeds are approximately 200 to date. Taranto seas, Mar Grande and Mar Piccolo, host 43 NIS of which 15 are macroalgal species; their presence is related to the pathways of introduction which typify the site (Cecere et al., 2016). Here we further report the accidental observation of two NIS considered among the worst in terms of negative impact on biodiversity and ecosystem functioning: *Caulerpa cylindracea* Sonder and *Asparagopsis* sp. *Caulerpa cylindracea* (as *Caulerpa racemosa* (Forsskål) J. Agardh var. *cylindracea* (Sonder) Verlaque, Huisman et Boudouresque) was firstly recorded along the coasts of North Africa in the 80's; the presence of this seaweed in the Gulf of Taranto dates back to 1996. The thalli consist in photosynthetic sub-cylindrical/claviform ramules attached to stolons. After a rapid colonization of the Taranto marine area, due to its impressive invasive ability (stolon elongation approx. 20mm d⁻¹), a regression of coverage was observed almost to its disappearance. In May 2022, only one erect axe was again observed in the Mar Piccolo of Taranto, at 1 meter of depth. In July 2022, a scattered but dense coverage of *C. cylindracea* was observed colonizing rocks in the intertidal zone, few centimetres of depth in the Gulf of Taranto; at the same site, in September it seemed to be disappeared. In June and July 2022, thalli of *Asparagopsis* sp. were observed on rocky bottom in the Mar Grande of Taranto, at 1 meter of depth. The genus *Asparagopsis* comprehends two species *A. armata* and *A. taxiformis* featuring a triphasic heteromorphic cycle. The gametophytic stage is represented by pink thalli attached to the substratum with pyramidal structure, and feathery branches; *A. armata* differs for the presence of harpoon-like branchlets. The tetrasporophyte stage appears as filamentous turfs and was previously considered as a separated species, i.e. *Falkenbergia hillebrandii* (Bornet) Falkenberg. Moreover, while *A. armata* is considered genetically uniform, for *A. taxiformis* molecular analysis showed the existence of 4 different evolutionary lines, two of them present in the Mediterranean Sea (Zanotta et al., 2014). In 1992, the presence of *A. armata* on the bottom of Cheradi Islands in front of the city of Taranto was reported. The allochthonous lineage of *A. taxiformis* was probably observed in 2015 in the Mar Piccolo of Taranto (Bottalico et al., 2015). The next step includes the use of molecular analysis on macroalgal samples to correctly identify the species.

Letteratura citata

- Bottalico A., Russo C., Pati R. (2015). Sulla presenza del genere *Asparagopsis* Montagne (Bonnemaisonales, Rhodophyta) in Puglia (Italia meridionale). Contributi scientifici Sezioni Regionali e Gruppi S.B.I. pp. 296-298 in Informatore Botanico Italiano, 47(2): 291-313.
- Cecere E., Petrocelli A., Belmonte M., Portacci G., Rubino F. (2016). Activities and vectors responsible for the biological pollution in the Taranto Seas (Mediterranean Sea, southern Italy): a review. Environmental Science and Pollution Research, 23: 12797-12810. DOI: 10.1007/s11356-015-5056-8
- Tsirintanis K., Azzurro E., Crocetta F., Dimiza M., Froglio C., Gerovasileiou V., Langeneck J., Mancinelli G., Rosso A., Stern N., Triantaphyllou M., Tsiamis K., Turon X., Verlaque M., Zenetos A. and Katsanevakis S. (2022). Bioinvasion impacts on biodiversity,



ecosystem services, and human health in the Mediterranean Sea. *Aquatic Invasion*, 17(3): 308-352.
<https://doi.org/10.3391/ai.2022.17.3.01>

Zanotta M., Carmona R., De La Rosa J., Salvador N., Sherwood A.R., Andreakis N., Altamirano M. (2014). Morphological differentiation of cryptic lineages within the invasive genus *Asparagopsis* (Bonnemaisoniales, Rhodophyta). *Phycologia*, 53: 233-242.
<https://doi.org/10.2216/13-247.1>

AUTORI

Giuseppe Denti (giuseppe.denti@irsa.cnr.it; Autore di riferimento), Antonella Petrocelli, Ester Cecere, Giovanni Fanelli, Loredana Papa, Fernando Rubino, CNR-Istituto di Ricerca sulle Acque (IRSA), sede di Taranto Talassografico "A. Cerruti", Via Roma 3, 74123, Taranto, Italia.



Allelopathic effects in the invasive macroalga *Rugulopterix okamurae* (Phaeophyceae): potential role of the newly discovered compound Dilkamural

D. Lenzo, A. Bartual, B. Úbeda, E. Zubía, B. Cuevas, M. A. Colangelo, A. Pasteris, L. Pezzolesi

Climate change is altering the distribution of organisms across the globe and is likely to change co-occurrence patterns and interspecific interactions of native, and invasive species, whose negative impact on marine ecosystems could affect the biodiversity and economy of coastal areas (Salvaterra et al., 2013). In aquatic environments, macroalgae provide habitat for fauna but are also able to produce different allelopathic compounds with high structural variability, that play an important role in species' successions (Pezzolesi et al., 2021). Whether invasive algae could produce allelopathic and toxic compounds, and how they could potentially change the population dynamics of native species remained unknown. In the last decade, the brown macroalga *Rugulopterix okamurae* has shown an intensive proliferation in the southwestern coasts of Europe (Strait of Gibraltar). Currently, there is no clue regarding the reasons of such huge invasive potential, although the involvement of chemical defenses has recently been suggested. Chemical study on *R. okamurae* from the Strait of Gibraltar led to the isolation of different secondary metabolites, among which the compound Dilkamural stands out because of its high concentration (Casal-Porras et al., 2021). In this context, this work was aimed to investigate the effects of Dilkamural on unicellular phototrophs to understand its potential role as allelochemical. In particular, this ecotoxicological study was carried testing two different species of microalgae (the diatoms *Phaeodactylum tricornutum* and *Cyclotella cryptica*) and a cyanobacterium (*Synechococcus elongatus*, PCC 7002) with different concentrations of Dilkamural. The effects on cells were evaluated for each species after 0, 3, 24, 48, 72 hours, adding "SYTOX green" to mark dead cells and analysing them through image flow cytometry (IFC) (Amnis ImageStream XMkII, Luminex Corporation). Results showed an acute effect of Dilkamural, after only three hours, on all the analysed species; particularly, effects on the viability and cell morphology of *P. tricornutum*, as well as cell integrity and reproduction of *S. elongatus* and *C. cryptica* were reported. The observed effects highlighted as the production of Dilkamural by this invasive species could affect algal population dynamics, reporting allelopathic effects on target organisms, thus potentially reducing the biodiversity of the coastal ecosystems where this invasive algal species has been introduced and extensively proliferates.

Letteratura citata

- Salvaterra, T., Green, D.S., Crowe, T.P. (2013). Impacts of the invasive alga *Sargassum muticum* on ecosystem functioning and food web structure. *Biol Invasions* 15, 2563–2576.
- Pezzolesi L, Accoroni S, Rindi F, Samorì C, Totti C, Pistocchi R. (2021). Survey of the allelopathic potential of Mediterranean macroalgae: production of long-chain polyunsaturated aldehydes (PUAs). *Phytochemistry*. 10.1016.
- Casal-Porras, I., Zubía, E., Brun, G.F. (2021). Dilkamural: A novel chemical weapon involved in the invasive capacity of the alga *Rugulopterix okamurae* in the Strait of Gibraltar. *Estuarine, Coastal and Shelf Science*, Volume 257, 107398, ISSN 0272-7714.

AUTORI

Denise Lenzo, (denise.lenzo2@unibo.it; Autore di riferimento), Laura Pezzolesi, Andrea Pasteris, Maria Antonia Colangelo, Dipartimento di Scienze Biologiche, Geologiche e Ambientali (BiGeA), Università di Bologna, Via Sant'Alberto 163, Ravenna, Italia.

Ana Bartual, Barbara Úbeda, Instituto Universitario de Investigaciones Marinas (INMAR), Campus de Excelencia Internacional del Mar (CEI-MAR), Universidad de Cádiz, Puerto Real, 11510 (Cádiz), Spagna.

Eva Zubía, Belen Cuevas, Departamento de Química Orgánica, Facultad de Ciencias del Mar y Ambientales, Universidad de Cádiz, 11510-Puerto Real, (Cádiz), Spagna.



Study of the effects of metabolites of invasive *Caulerpa* spp. on *Posidonia oceanica* seedling growth: unexpected effects of the vehicle (DMSO)

D. Oliva, A. Piro, M. Carbone, E. Mollo, F. Scarcelli, V. Osso, D.M. Nisticò, S. Mazzuca

The Mediterranean hosts the largest number of allochthonous macrophytes (Zenetos et al., 2012) that can replace indigenous species, causing a decrease in the biodiversity of native species and altering the ecosystem. The species of greatest concern are mainly *Caulerpa taxifolia* and *Caulerpa cylindracea*, both infesting tropical algae whose role towards the biodiversity of Mediterranean macrophytes has not yet been clarified (Marbà et al., 2005). The algae of the genus *Caulerpa* also have the ability to compete with seagrasses (Williams SL 2007). It has been hypothesized that, at the basis of this competition, there may be an allelopathic interaction between species (Defranoux et al, 2021). Among the secondary metabolites produced by algae, the alkaloid *caulerpin* and the sesquiterpene *caulerpenyne* have been considered to play a role in the regression of endemic communities (Piazzi et al., 2001); their mechanism of action, however, is currently unknown.

The aim of the present research is to clarify the effects of molecules purified from *C. cylindracea* on growth, phenology, and molecular responses of *Posidonia oceanica* cuttings acclimated under controlled conditions in a mesocosm, and thus identify the metabolic pathways that are induced by the chemical interaction between the species. Here we will present the *i*) optimized conditions for the cultivation of *P. oceanica* cuttings in the aquarium, *ii*) the results of the extraction, purification and characterization of the main secondary metabolites in *Caulerpa* spp. and *iii*) the results of a pilot experiment on *P. oceanica* treated with *caulerpin* and *caulerpenyne* dissolved in DMSO (dimethyl sulfoxide). An unexpected result will also be presented which concerns the effect of DMSO on the studied parameters. This data will be discussed in relation to the recent publication which has shown that, among the seagrass, *P. oceanica* is the largest producer of DMSO together with its precursor, dimethylsulfoniopropionate (DMSP) (Richir et al., 2021).

Letteratura citata

- Marbà, N., Duarte, C.M., Díaz-Almela, E. et al. (2005) Direct evidence of imbalanced seagrass (*Posidonia oceanica*) shoot population dynamics in the Spanish Mediterranean. *Estuaries* 28, 53–62
- Williams, S. L. (2007) Introduced species in seagrass ecosystems: status and concerns. *Journal of Experimental Marine Biology and Ecology*, 350(1-2), 89-110.
- Defranoux F., Noè S., Cutignano A., Casapullo A., Ciavatta M.L., Carbone M., et al. (2021) Chemoecological study of the invasive alga *Caulerpa taxifolia* var. *distichophylla* from the Sicilian coast. *AQUATIC ECOLOGY*.
- Piazzi, L., Ceccherelli, G., & Cinelli, F. (2021) Threat to macroalgal diversity: effects of the introduced green alga *Caulerpa racemosa* in the Mediterranean. *Marine Ecology Progress Series*, 210, 149–159.
- Jonathan, Richir & Champenois, W. & de Fouw, Jimmy & Borges, Alberto. (2021) Dimethylsulfoniopropionate and dimethylsulfoxide in *Posidonia oceanica*. *Marine Biology*. 168. 10.1007/s00227-021-03961-5.

AUTORI

Daniela Oliva (daniela.oliva@unical.it; Autore di riferimento), Amalia Piro (amalia.piro@unical.it), Lab. Biologia e Proteomica Vegetale (Lab.Bio.Pro.Ve.), Dip C.T.C., Università della Calabria, Rende, Italia.
 Marianna Carbone (mcarbone@icb.cnr.it), Ernesto Mollo (emollo@icb.cnr.it), Consiglio Nazionale delle Ricerche (CNR), Institute of Biomolecular Chemistry (ICB), Via Campi Flegrei, 34, 80078 Pozzuoli (Na), Italia.
 Faustino Scarcelli (faustino.scarcelli@unical.it), Vincenzo Osso (vincenzo.osso@unical.it), Dante Matteo Nisticò (dante.nisticò@unical.it), Silvia Mazzuca (silvia.mazzuca@unical.it), Lab. Biologia e Proteomica Vegetale (Lab.Bio.Pro.Ve.), Dip C.T.C., Università della Calabria, Rende, Italia.



Algal assemblages' composition of two semi-submerged marine caves at Tremiti Islands Marine Protected Area (Italy, Adriatic Sea)

A. Tursi, F. Mastrototaro, G. Chimienti, A. Bottalico

Semi-submerged marine caves are peculiar coastal habitats widely spread all over the Mediterranean Sea (Gerovasileiou & Bianchi, 2021). These environments enrich the coastal landscape complexity, being extremely important for marine biodiversity. In fact, they represent valuable hotspots of biodiversity, being colonized by peculiar species generally found at higher depths, well adapted to semi-submerged marine caves characterized by low light intensity and scarce hydrodynamism (Ouerghi et al., 2019). The benthic communities in these caves include a rich and diverse macroalgal flora which plays a key role in the stability of these habitats. Nevertheless, only a few studies have been carried out about the algal assemblages of such environments (Alongi et al., 2012; Bottalico et al., 2016). In this context, Tremiti Islands Marine Protected Area (MPA) hosts several semi-submerged marine caves, whose algal composition is scarcely known (Pignatti et al., 1967; Rizzi Longo, 1972). This study is focused on the monitoring of two semi-submerged marine caves in the MPA, namely Viole and Rondinelle caves. A qualitative sampling was carried out by SCUBA diving in two different monitoring periods (spring and autumn 2021). A total of 42 species were identified in the two semi-submerged marine caves. At Viole cave, Rhodophyta showed to be the most abundant taxa with 24 species, followed by Chlorophyta and Ochrophyta-Phaeophyceae, which were represented by 7 and 6 species, respectively. A total of 20 algal species was observed at Rondinelle Cave, with 13 Rhodophyta, 5 Chlorophyta and 2 Ochrophyta-Phaeophyceae. In both caves, Rhodophyta were particularly abundant in the inner portions of the caves, where the light intensity was scarce, while Chlorophyta and Ochrophyta-Phaeophyceae tended to be more abundant nearby the entrance and the end of the caves, which had collapsed vaults, allowing the presence of sunlight. Cave walls were dominated by encrusting algae belonging to the genus *Peyssonnelia*, confirming what observed more than 50 years earlier (Pignatti et al., 1967). This study contributed to implementing the knowledge about the algal assemblages of two semi-submerged marine caves, particularly important for the tourism industry of Tremiti Archipelago. Our results showed the presence of several algal species of great ecological and conservation values, highlighting the importance of monitoring and conservation of these coastal habitats. Future perspectives will focus on the algal communities of other semi-submerged marine caves of the MPA, in order to identify any similarities and differences, possibly also related to different environmental conditions and/or to anthropic actions.

Letteratura citata

- Alongi G., Cormaci M., Furnari G., Catra M. (2012). Floristic macroalgal diversity in selected submarine caves located within two marine protected areas off Lampedusa Island and Sicily (Italy). *Botanica Marina*, 55(4): 387–397.
- Bottalico A., Alongi G., Perrone C. (2016). Macroalgal diversity of Santa Cesarea-Castro (Salento Peninsula, southeastern Italy). In *Anales del Jardín Botánico de Madrid*, 73(2): 1–12.
- Gerovasileiou V., Bianchi C. N. (2021). Mediterranean marine caves: A synthesis of current knowledge. *Oceanography and Marine Biology*, 1–87.
- Rizzi Longo L. (1972). La flora sottomarina delle Isole Tremiti. *Istituto Veneto di Scienze, Lettere ed Arti*, 130:329–376.
- Ouerghi A., Gerovasileiou V., Bianchi C.N. (2019). Mediterranean marine caves: A synthesis of current knowledge and the Mediterranean Action Plan for the conservation of 'dark habitats'. In *Marine Caves of the Eastern Mediterranean Sea*, 53: 1–13.
- Pignatti S., De Cristini P., Rizzi L. (1967). Le associazioni algali della grotta delle Viole nell'isola di S. Domino (Is. Tremiti). *Plant Biosystem*, 101(2): 117–126.

AUTORI



Andrea Tursi (andrea.tursi@uniba.it; Autore di riferimento), Francesco Mastrototaro (francesco.mastrototaro@uniba.it), Giovanni Chimienti (giovanni.chimienti@uniba.it), Antonella Bottalico (antonella.bottalico@uniba.it), Department of Bioscience, Biotechnology and Environment, University of Bari Aldo Moro, Via Orabona 4, 70124 Bari, Italy.



New insights into the reproductive strategies of the endangered *Fucus virsoides* useful for implementing restoration measures

S. Kaleb, E. Descourvières, V. Bandelj, F. Gianni, M. Orlando-Bonaca, A. Falace

Canopy-forming algae of the order Fucales and Laminariales (Phaeophyceae) are among the most valuable marine species in temperate waters. They provide a number of important ecosystem services, including carbon sequestration, shelter and nursery (Fabbrizzi et al., 2020). They also play a central role in understory growth by influencing light, desiccation and water movement. *Fucus virsoides* J. Agardh (Fucales, Phaeophyta), endemic to the Adriatic Sea (Linardić, 1949; Giaccone & Pignatti, 1967; Munda, 1972), is a glacial relict that has survived in the Adriatic Sea thanks to the specificities of this basin (Munda, 1972, Orlando-Bonaca et al., 2013). In the past, its range extended from the Venice Lagoon to the Boka Kotorska (Mačić, 2006). In recent decades, it has experienced dramatic declines or local extinctions (e.g. Battelli, 2016; Mačić, 2006; Orlando-Bonaca et al., 2013). Due to its endangered status and ecological role, it has been included in the "List of Threatened or Endangered Species" of the Barcelona Convention (amended Annex II of the Protocol on Specially Protected Areas and Biological Diversity in the Mediterranean, UNEP; 2019).

The lack of adult plants producing recruits hinders the natural recovery of degraded areas. Therefore, active restoration is recommended to recover populations within a reasonable time frame. In this study, for the first time, an ex situ pilot restoration of *F. virsoides* was carried out by outplanting cultivated recruits in the Gulf of Trieste. The following aspects were addressed: the efficiency of seedling culture in the laboratory; the effectiveness of an intermediate step in the field in suspended structures; and the subsequent anchoring on rocky substrate. Fertile apical fronds were collected from the donor site and the development of the recruits on clay tiles was followed under controlled conditions. Two different enriched culture media were tested. The recruits were further cultivated on lantern nets at the recipient site until they reached a "refuge size".

The effectiveness of restoration was measured by quantifying juvenile cover, growth and survival at different time points. This study enlarges the biological knowledge of the early life and juvenile stages of *F. virsoides* and draws some practical methodological considerations for implementing restoration measures. It provides new information to optimise best practises in laboratory culture and addresses the first steps of recovery and introduction of juvenile thalli into the natural environment.

Letteratura citata

- Battelli C. (2016). Disappearance of *Fucus virsoides* J. Agardh from the slovenian coast (Gulf of Trieste, Northern Adriatic). Annals for Istrian and Mediterranean Studies. 26: 1-12. doi.org/10.19233/ASHN.2016.1.
- Fabbrizzi E., Scardi M., Ballesteros E., Benedetti-Cecchi L., Cebrian E., Ceccherelli G., De Leo F., Deidun A., Guarnieri G., Falace A., Fraissinet S., Giommi C., Mačić V., Mangialajo L., Mannino A.M., Piazzi L., Ramdani M., Rilov G., Rindi L., Rizzo L., Sarà G., Souissi J.B., Taskin E., Frachetti S. (2020). Modeling Macroalgal Forest Distribution at Mediterranean Scale: Present Status, Drivers of Changes and Insights for Conservation and Management. Front. Mar. Sci. 7: 1-18. https://doi.org/10.3389/fmars.2020.00020.
- Giaccone G., Pignatti S. (1967). Studi sulla produttività primaria del fitobentos nel Golfo di Trieste. - II. La vegetazione del Golfo di Trieste. Nova thalassia. 3(2): 1-28.
- Linardić J. (1949). Studije o jadranskem fukusu (*Fucus virsoides*). Acta Botanica. 12, 13: 7-131.
- Mačić V. (2006). Distribution of seaweed *Fucus virsoides* J. Agardh in Boka Kotorska Bay (South Adriatic Sea). Annales, Ser. Hist. Nat., Koper. 16: 1-4.
- Munda M.I. (1972). Seasonal and ecologically conditioned variations in the *Fucus virsoides* association from the Istrian coast (Northern Adriatic). Dissertationes Slovenska Akademija Znanosti in Umetnosti. 15: 1-33.
- Orlando-Bonaca M., Mannoni P.A., Poloniato D., Falace A. (2013). Assessment of *Fucus virsoides* distribution in the Gulf of Trieste (Adriatic Sea) and its relation to environmental variables. Botanica Marina. (56)5-6: 451-459. <https://doi.org/10.1515/bot-2013-0027>



AUTORI

Sara Kaleb, Università degli Studi di Trieste, Via Licio Giorgieri 10, 34127 Trieste.
Emanuelle Descourvières, Università degli Studi di Trieste, Via Licio Giorgieri 10, 34127 Trieste; Istituto Nazionale di Oceanografia e di Geofisica Sperimentale - OGS, Via Beirut 2, 34151 Trieste.
Vinko Bandelj, Istituto Nazionale di Oceanografia e di Geofisica Sperimentale - OGS, Via Beirut 2, 34151 Trieste.
Fabrizio Gianni, Istituto Nazionale di Oceanografia e di Geofisica Sperimentale - OGS, Via Beirut 2, 34151 Trieste.
Martina Orlando-Bonaca, National Institute of Biology (NIB), Marine Biology Station Piran, Fornače 41, 6330 Piran, Slovenia.
Annalisa Falace (falace@units.it; Autore di riferimento), Università degli Studi di Trieste, Via Licio Giorgieri 10, 34127 Trieste.



Sessione Microalge II

Moderatore: Stefano Accoroni



Search of freshwater algal specie with adaptive mechanisms to Cr (VI) stress

M. Marieschi, G. Molinari, S. Savi, R. Bolpagni, A. Torelli

Sulfate uptake/assimilation has a pivotal role in enhancing cell defense and on the onset of tolerance mechanisms toward abiotic stresses, in particular to heavy metals, chromium included. This phenomenon, known as SIR/SED (Sulfur Induced Resistance/Sulfur Enhanced Defense) (Rausch and Wachter 2005), relies on a high capability to synthesize cysteine (cys) and reduced glutathione (GSH). These molecules are directly involved in chelating various metal ions, Cr(VI) included, and in the reduction of metal induced oxidative stress. In previous studies we observed that differences in sulfate pathway are involved in different Cr(VI) sensitivity in two strains of *Scenedesmus acutus* (Gorbi et al 2007, Marieschi et al. 2015, Sardella et al 2019, Ferrari et al 2022). To verify if an enhanced cys production could represent a mechanism of environmental adaptation, we tried to isolate microalgae in habitats subject to abiotic stress to be addressed to laboratory research. Algal samples were collected in a chromigen spring at Mount Prinzera, an ophyolithic mountain nearby Parma. The spring water shows Cr(VI) levels above low limits, with an average of 12 Cr(VI) µg/l depending on water flow. The eco-toxicological assay for algal growth (OECD 201/2006) indicate that the water has an inhibitory effect on algal growth. This sampling area thus proved to be an optimal source for the isolation of organisms which could have evolved tolerance mechanisms to Cr(VI). We isolated nearly 60 colonies, among them some of green algae, diatoms and cyanobacteria. Isolation of pure colonies and species identification is still in progress. Some of the tested species shows a Cr(VI) tolerance to concentration 10fold higher than of the spring in which they were isolated, thus constituting a good experimental material for future researches.

Letteratura citata

- Ferrari, M., Cozza, R., Marieschi, M. and Torelli, A. (2022). Role of Sulfate Transporters in Chromium Tolerance in *Scenedesmus acutus* M. (Sphaeropleales). Plants (Basel, Switzerland), [online] 11(2), p.223. doi:10.3390/plants11020223.
- Gorbi, G., Zanni, C. and Corradi, M.G. (2007). Sulfur starvation and chromium tolerance in *Scenedesmus acutus*: a possible link between metal tolerance and the regulation of sulfur uptake/assimilation processes. Aquatic Toxicology (Amsterdam, Netherlands), [online] 84(4), pp.457–464. doi:10.1016/j.aquatox.2007.07.006.
- Marieschi, M., Gorbi, G., Zanni, C., Sardella, A. and Torelli, A. (2015). Increase of chromium tolerance in *Scenedesmus acutus* after sulfur starvation: Chromium uptake and compartmentalization in two strains with different sensitivities to Cr(VI). Aquatic Toxicology, 167, pp.124–133. doi:10.1016/j.aquatox.2015.08.001.
- Rausch, T., Wachter, A., 2005. Sulfur metabolism: a versatile platform for launching defence operations. Trends in Plant Sci. 10, 503-509.
- Sardella A., Marieschi M., Mercatali I., Zanni C., Gorbi G. (2019). The relationship between sulfur metabolism and tolerance of hexavalent chromium in *Scenedesmus acutus* (Sphaeropleales): Role of ATP sulfurylase. Aquatic Toxicology, 216: 105320

AUTORI

Matteo Marieschi (matteo.marieschi@unipr.it), Giuseppe Molinari, Simone Savi, Rossano Bolpagni, Anna Torelli (anna.torelli@unipr.it; Autore di riferimento), Dipartimento di Chimica, Scienze della Vita e della Sostenibilità Ambientale, Università di Parma, Viale delle Scienze 11/A, 43124 Parma, Italia.



Interaction between OAS-TL and desulphydrase activity in Cr(VI) tolerance in the freshwater green alga *Scenedesmus acutus*

G. Molinari, M. Marieschi, R. Bolpagni, A. Torelli

In plant organisms, sulfur-mediated defense (SED) is crucial for limiting damage caused by heavy metals. Previous works, conducted on two strains of *Scenedesmus acutus*, Wt and chromium-tolerant (CrT), evidenced a transient tolerance increase following a period of sulfur starvation, related to the increase of sulfur uptake and assimilation, in both strains but significantly higher in the CrT strain leading to a higher cysteine content in this latter (Sardella et al. 2019).

Cysteine is synthesized by the enzymatic cysteine synthase complex (CSC), composed by a hexamer of Serine Acetyltransferase (SAT) and two dimers O-acetyl-Serine (Thiol) Lyase (OAS-TL) (Wirtz et al. 2006), present in different cell compartments with different role in cys homeostasis.

OAS-TL activity measured during Cr(VI) exposure preceded by 7-day preculture in standard (S-sufficient) or sulfur-deprived medium (S-replete), was strongly activated in both strains after S-starvation and dropped slowly in S-replete CrT strain exposed to chromium than in controls or in the Cr exposed Wt strain.

The significantly higher cysteine amount observed in the CrT strain during recovery after starvation was hypothesized to involve cytoplasmic rather than chloroplastic CSC complex (Sardella et al. 2019). This hypothesis poses however the question of sulfide production, since sulfite reductase (SIR) have apparent exclusively plastid localization. Extra plastidic sulfide production can be substantiated by OAS-TLs desulphydrase side activity (Riemenschneider et al. 2005) and from the activity of specific DES enzymes which can contribute to recycling organic sulfur. To verify possible differences between strains we measured L-Cys desulphydrase activity in the same conditions in which OAS-TL activity was assayed. Also in this case a 7 day S-starvation induced the increase of desulphydrase activity in both strain and a significantly slower decrease in the CrT strain exposed to 2 mg Cr(VI)/L. This trend clearly followed that of OAS-TL activity apparently confirming desulphydrase as OAS-TL side activity.

We further isolated a *S. acutus* sequence homologous to Arabidopsis D-Cys DES and analyzed its expression in response to S-starvation to verify if a different expression allowed to hypothesize a different sulfide production in the two strains. In Wt D-Cys DES is expressed at low levels in standard condition and is induced by S-deprivation, whereas in CrT DES levels are significantly higher than in Wt in standard medium and remains nearly stable in S-deprived medium.

A different relationship between D-Cys-DES and OAS-TL side activity in the two strains could underlie their different cysteine homeostasis and Cr(VI) tolerance. D-Cys DES activity remains however to be investigate to get more insight in extraplastidic sulfide production.

Letteratura citata

- Sardella A., Marieschi M., Mercatali I., Zanni C., Gorbi G. (2019). The relationship between sulfur metabolism and tolerance of hexavalent chromium in *Scenedesmus acutus* (Sphaeropleales): Role of ATP sulfurylase. Aquatic Toxicology, 216: 105320
 Wirtz M., Rüdiger H. (2006). Functional analysis of the cysteine synthase protein complex from plants: Structural, biochemical and regulatory properties. Journal of Plant Physiology, 163: 273-286
 Riemenschneider, A., Riedel, K., Hoefgen, R., Papenbrock, J. and Hesse, H. (2005). Impact of Reduced O-Acetylserine(thiol)lyase Isoform Contents on Potato Plant Metabolism. Plant Physiology, 137(3), pp.892-900. doi:10.1104/pp.104.057125.

AUTORI

Giuseppe Molinari (giuseppe.molinari@unipr.it), Matteo Marieschi, Rossano Bolpagni, Anna Torelli (anna.torelli@unipr.it; Autore di riferimento), Dipartimento di Chimica, Scienze della Vita e della Sostenibilità Ambientale, Università di Parma, Viale delle Scienze 11/A, 43124 Parma, Italia.



Effetti di composti fenolici presenti in Acque di Vegetazione sulle microalghe

L. Mollo, F. Drigo, A. Norici

La fitodepurazione di acque reflue attraverso microalghe è un processo biotecnologico che, attraverso il metabolismo mixotrofo di alcune specie algali e il loro elevato tasso di crescita, permette di abbassare i costi della coltivazione industriale di microalghe e, al contempo, depurare uno scarto potenzialmente inquinante.

Fra le varie tipologie di acque reflue di potenziale interesse per la coltivazione di microalghe vi sono le Acque di Vegetazione (AV), un sottoprodotto potenzialmente tossico (Sakarita et al., 2020) derivante dall'estrazione dell'olio d'oliva. Queste acque di scarto sono ricche in elementi minerali e materia organica (Komnitsas et al., 2012), ma la presenza di composti fenolici ne limita il loro utilizzo come mezzo di crescita per microalghe.

Le AV possono infatti contenere fino 1-2 g L⁻¹ (Deeb et al., 2012) di composti fenolici che, se rilasciati in modo incontrollato nell'ambiente, possono causare effetti dannosi per le comunità microbiche del suolo e delle acque (Bouarab-Chibane et al., 2019); una loro rimozione è quindi necessaria. Nel caso in cui si voglia percorrere la via sostenibile della fitodepurazione mediante microalghe, è necessario comprendere appieno l'effetto dei composti fenolici su questi organismi. Ad oggi, gli effetti dei fenoli delle AV sulle microalghe non sono del tutto noti (Lindner & Pleissner, 2019) e studi più approfonditi sono stati svolti solamente per composti fenolici di origine antropica e farmaceutica (Surkatt & Al-Zuhair, 2018).

In quest'ottica 10 diverse microalghe, scelte sulla base della loro origine filogenetica e presenti sia in mare che in acque dolci, sono state esposte a tre fenoli (tirosolo, acido cumarico e acido caffeoico) tipici delle AV. Sono stati quindi studiati i tassi di crescita e le rimozioni percentuali dei composti fenolici e si è valutata la presenza di risposte comuni alle specie in esame. Come già osservato, le risposte si sono rivelate specie-specifiche; tuttavia, è stata osservata una maggiore tolleranza, in termini di crescita, delle specie d'acqua dolce rispetto a quelle marine. Non è stata riscontrata una correlazione fra crescita e rimozione di composti fenolici, equivalente fra le due categorie, suggerendo quindi l'esistenza di meccanismi passivi di adsorbimento sulla parete cellulare (Wu et al., 2022).

Gli effetti citotossici e ossidativi dei fenoli sono stati inoltre studiati su *Dunaliella salina*, unica microalga marina in esame capace di sopravvivere ai tre composti fenolici: indice di riduzione, DPPH assay e Comet assay sono stati impiegati per studiare gli effetti dei fenoli sullo stato di ossidazione della cellula e a livello di DNA.

Ciò che è emerso è la comparsa di risposte rapide in seguito all'esposizione ai fenoli con successiva acclimatazione e riduzione della risposta nel tempo.

La comprensione degli effetti a livello cellulare dei composti fenolici apre le porte a una fitodepurazione più efficiente e a un utilizzo più consci delle microalghe in questi processi biotecnologici.

Letteratura citata

- Sakarika, M.; Koutra, E.; Tsafarakidou, P.; Terpou, A.; Kornaros, M. Microalgae-Based Remediation of Wastewaters. In Microalgae Cultivation for Biofuels Production; Elsevier, 2020; pp. 317–335 ISBN 978-0-12-817536-1.
- Komnitsas, K.; Zaharaki, D. Pre-Treatment of Olive Mill Wastewaters at Laboratory and Mill Scale and Subsequent Use in Agriculture: Legislative Framework and Proposed Soil Quality Indicators. Resources, Conservation and Recycling 2012, 69, 82–89, doi:10.1016/j.resconrec.2012.09.009.
- Deeb, A.A.; Fayyad, M.K.; Alawi, M.A. Separation of Polyphenols from Jordanian Olive Oil Mill Wastewater. Chromatography Research International 2012, 2012, 1–8, doi:10.1155/2012/812127.
- Bouarab-Chibane, L.; Forquet, V.; Lantéri, P.; Clément, Y.; Léonard-Akkari, L.; Oulahal, N.; Degraeve, P.; Bordes, C. Antibacterial Properties of Polyphenols: Characterization and QSAR (Quantitative Structure–Activity Relationship) Models. Front. Microbiol. 2019, 10, 829, doi:10.3389/fmicb.2019.00829.
- Lindner, A.V.; Pleissner, D. Utilization of Phenolic Compounds by Microalgae. Algal Research 2019, 42, 101602, doi:10.1016/j.algal.2019.101602.



Surkatt, R.; Al-Zuhair, S. Microalgae Cultivation for Phenolic Compounds Removal. Environ Sci Pollut Res 2018, 25, 33936–33956, doi:10.1007/s11356-018-3450-8.

Wu, P.; Zhang, Z.; Luo, Y.; Bai, Y.; Fan, J. Bioremediation of Phenolic Pollutants by Algae - Current Status and Challenges. Bioresource Technology 2022, 350, 126930, doi:10.1016/j.biortech.2022.126930.

AUTORI

Lorenzo Mollo (l.mollo@pm.univpm.it; Autore di riferimento), Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona, Italy

Filippo Drigo (S1099217@studenti.univpm.it), Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona, Italy

Alessandra Norici (a.norici@univpm.it), Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona, Italy



Benthic diatom communities on seagrass and corals in Arabian Gulf: floristic analysis, population dynamics and estimate of contribute to the primary production of the basin

C. Auciello, C. Pennesi, C. Ciniglia, M. Iovinella, L. Rabaoui, M. De Stefano

Primary productivity generated by phanerogams, macroalgae, phytoplankton and microphytobenthos account for over 40% of entire world productivity (Duarte & Cebrih, 1996). Phytoplankton is considered the main global producer whereas macroalgae, distributed approximately for 1.8% of marine area, exhibit estimated values of primary productivity of 2.55×10^9 tC/y (Charpy and Sournia, 1990). Primary productivity attributed to microphytobenthos (with an estimated distribution similar to those attributed to macroalgae) not exceed 0.34×10^9 tC/y (Charpy and Sournia, 1990) this demonstrating a strong underestimation of the ecological contribution of microphytobenthic communities if we considering their wide distribution on all organic and inorganic substrates (Fe, 1971). All this is even truer considering that main composer of microphytobenthic communities are generally diatoms (Evrard, et al., 2012), microalgae characterized by a high photosynthetic efficiency and high rate of primary productivity (Falkowski & Raven, 1997). Therefore, we can assume that in order to study the contribution to net primary productivity of microphytobenthos in a basin, we could mainly study the primary productivity of its diatom communities. Our study, entirely based on scanning electron microscopy, aimed to decrypt and document the hidden biodiversity of diatoms associated to seagrasses and hard corals along the Saudi coasts of Arabian Gulf and to estimate their real contribute to the primary productivity of basin. Preliminary results demonstrated that both seagrasses and corals hosted benthic diatom communities who constituted the major element of their epibenthic microalgal flora. Significant differences in terms of cell density, species composition and community structure were observed between the diatom communities of seagrasses and corals and within the different sampling locations. In particular, cell densities of seagrass diatom communities are more than one hundred times higher than those of corals with mean values approximately of 6000-7000 cell/mm² in seagrass vs 100-300 cell/mm² in corals. Within the community, *Cocconeis* was the dominant genus represented by four species. The abundances are higher in the central stations of the gulf than in the north and south stations, most likely due to the particular environmental conditions that exist in these two areas of the gulf. The estimated primary productivity of the epiphytic diatom communities within the Arabian Gulf account for 24×10^6 tC/y.

Letteratura citata

- Charpy-Roubaud, C., A. Sournia. (1990). The comparative estimation of phytoplanktonic, microphytobenthic and macrophytobenthic primary production in the oceans. Mar. Micr. Food Webs. 4: 31-57.
 Duarte, C. A., Cebrih, J. (1996). The fate of marine autotrophic production. Limnol. Oceanogr. 41(8): 1758-1766.
 Evrard, V., Huettel, M., Cook, P., Soetaert, K., Heip, C.H.R., Middelburg, J.J. (2012). Importance of phytodetritus and microphytobenthos for heterotrophs in a shallow subtidal sandy sediment. Mar. Ecol. Prog. Ser. 455: 13-31.
 Falkowski P.G. & Raven J.A. (1997). Aquatic photosynthesis. Blackwell, Oxford. 375.
 Fe, R. (1971). Benthic marine diatoms. Oceanogr. Mar. Biol. Annu. 9: 83-139.

AUTORI

Concetta Auciello (concetta.auciello@unicampania.it; Autore di riferimento), Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "L. Vanvitelli", via Vivaldi 43, 81100 Caserta, Italia.
 Chiara Pennesi (chiara.penesi@szn.it), Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn, C.da Torre Spaccata, Località Torre Spaccata 87071 Amendolara (CS), Italia.
 Claudia Ciniglia (claudia.ciniglia@unicampania.it), Manuela Iovinella (manuela.iovinella@unicampania.it), Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "L. Vanvitelli", via Vivaldi 43, 81100 Caserta, Italia.



Lotfi Rabaoui (lrabaoui@gmail.com), King Fahd University of Petroleum and Minerals, Dhahran, Eastern Province, Saudi Arabia.

Mario De Stefano (mario.destefano@unicampania.it), Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "L. Vanvitelli", via Vivaldi 43, 81100 Caserta, Italia.



Potential of *Nannochloropsis oceanica* for bioremediation

M. Blasio, G.V. de Klerk, P. Marotta, A. Cutignano, G. Trancone, G. Dupont, A. Sardo, P. De Luca, S. Papirio, S Balzano

Heavy Metals (HMs) are widespread in nature at trace concentrations, nevertheless, anthropogenic activities, including mining, agriculture, and manufacturing, cause an excessive discharge of HMs in the environment. Conventional technologies include chemical precipitation, electrochemical techniques, ion exchange, membrane filtration, coagulation and flocculation. In recent years, HM removal from wastewaters using living or dead organisms, including microalgae, has emerged as a valid alternative to conventional methods (Manikandan et al., 2022). The small size of unicellular organisms, including microalgae, leads to a great surface/volume ratio and a higher availability of HM binding sites on their cell surface compared to multicellular organisms (Cameron et al., 2018). In this study, we investigated the HM removal potential of *Nannochloropsis oceanica* CCMP1779, as well as two mutant strains engineered using CRISPR/Cas9 technology to knock out the polyketide synthase (PKS) gene. PKS is likely to be involved in the biosynthesis of the building blocks of the cell wall polymer algaenan. In detail, we evaluated the ability of *N. oceanica* wild-type and two mutant strains to grow for 10 days in the presence of chromium, copper, lead and zinc at concentrations similar to those observed in polluted hotspots of the Gulf of Naples. At late exponential phase, the biomass was harvested by centrifugation and HM concentrations were determined in both algal pellet and supernatant by inductively coupled plasma mass spectroscopy (ICP-MS). The results indicate that microalgal growth is poorly affected by the presence of HMs reaching a cellular concentration between 3 and 5×10^7 cell ml⁻¹. In addition, the ICP-MS analysis highlighted an impressive ability of *Nannochloropsis* cultures to remove HMs from the environment. The two mutant strains, KO9 and KO59, exhibited a higher cellular content of chromium, zinc, and lead compared to the wild-type; the mutant KO9 accumulated greater amounts of cadmium and copper compared to the wild type and the mutant KO59. A fundamental role in HM removal is covered by functional groups present on microalgal cell wall surface. A putative modification in cell wall chemical composition occurring in mutant strains may result in a higher affinity toward HMs. Current results suggest a great potential of *Nannochloropsis oceanica* for the HM removal from contaminated sea water. In particular, mutants with putative alteration in cell wall chemistry exhibited a higher performance compared to the wild-type.

Letteratura citata

- Cameron, H., Mata, M.T., and Riquelme, C. (2018) The effect of heavy metals on the viability of *Tetraselmis marina* AC16-MESO and an evaluation of the potential use of this microalga in bioremediation. *PeerJ*. 6.
- Díaz, Y.A., Martínez, E.M., García, L.C., Serrano, M.C., and Sierra, E.M. (2022) Phycoremediation as a Strategy for the Recovery of Marsh and Wetland with Potential in Colombia. *Resour 2022, Vol 11, Page 15.* 11: 15.
- Leong, Y.K., and Chang, J.S. (2020) Bioremediation of heavy metals using microalgae: Recent advances and mechanisms. *Bioresour Technol.* 303: 122886.
- Manikandan, A., Suresh Babu, P., Shyamalagowri, S., Kamaraj, M., Muthukumaran, P., and Aravind, J. (2022) Emerging role of microalgae in heavy metal bioremediation. *J Basic Microbiol.* 62: 330–347.

AUTORI

- Martina Blasio (martinablasio@szn.it; Autore di riferimento), Gioia Vanessa de Klerk (gioia.vanessa.deklerk@imbrsea.eu), Dipartimento di Biotecnologie Marine Ecosostenibili (BlueBio), Stazione Zoologica Anton Dohrn, Villa Comunale, 80133, Napoli, Italia.
- Pina Marotta (pina.marotta@szn.it) Dipartimento Infrastrutture di Ricerca per le Risorse Biologiche Marine (RIMAR), Stazione Zoologica Anton Dohrn, Villa Comunale, 80121, Napoli.
- Adele Cutignano (adele.cutignano@icb.cnr.it), Istituto di Chimica Biomolecolare (ICB), Consiglio Nazionale delle Ricerche (CNR), Via Campi Flegrei, 34, 80078, Pozzuoli (Na), Italia.



Gennaro Trancone (gennaro.trancone@unina.it), Department of Civil, Architectural, and Environmental Engineering, University of Naples Federico II, Via Claudio, 21, 80125, Napoli, Italia.

Guillaume Dupont (guillaume.dupont@imbrsea.eu), Angela Sardo (angela.sardo@szn.it), Dipartimento di Biotecnologie Marine Ecosostenibili (BlueBio), Stazione Zoologica Anton Dohrn, Villa Comunale, 80133, Napoli, Italia.

Pasquale De Luca (pasquale.deluca@szn.it), Dipartimento Infrastrutture di Ricerca per le Risorse Biologiche Marine (RIMAR), Stazione Zoologica Anton Dohrn, Villa Comunale, 80121, Napoli, Italia.

Stefano Papirio (stefano.papirio@unina.it), Department of Civil, Architectural, and Environmental Engineering, University of Naples Federico II, Via Claudio, 21, 80125, Napoli, Italia.

Sergio Balzano (sergio.balzanno@szn.it), Dipartimento di Biotecnologie Marine Ecosostenibili (BlueBio), Stazione Zoologica Anton Dohrn, Villa Comunale, 80133, Napoli, Italia.



A Photo Guide of Harmful Algae along Costal areas of Apulian Region

S. D'Arpa, M.R. Vadrucci, P.C. Ciciriello, E. Stanca A. Pastorelli, N. Ungaro

Harmful Algae Blooms (HABs) can cause serious negative impacts on the economy, fishery/seafood industry, environment, and even recreational activities. It should be stressed that some of these agents produce toxins that can cause human illness and even death. In particular, the presence of HABs species represent a great threat for bivalve aquaculture. The impact of their blooms is directly related to the harmfulness of the shellfish toxins they produce. Filter-feeding bivalves accumulate microalgal toxins, which can reach concentrations (regulatory levels) unsafe for human consumption and thus health and fisheries authorities have to implement management measures, including the closure of shellfish harvesting areas. The technical guidance for the application of EU Regulation identified the criteria for defining monitoring plans for the quality control of bivalve for human consumption and includes, among the parameters to be monitored, the presence of toxin-producer phytoplankton in the area of shellfish harvesting.

In the Apulia region a monitoring programme finalized to the assessment of the quality of marine water for bivalve aquaculture started at the beginning of 2021 (according to D.G.R n. 2014/2019).

The monitoring plans included the analysis of phytoplankton in a number of 40 sampling points among shellfish farms, relaying areas and natural beds of bivalves with a biweekly frequency.

Previous studies on HABs dynamics along the Apulian coasts and transitional waters showed the presence of phytoplankton toxin (ASP and DSP producers) and high-biomass producer (Roselli et al., submitted). Therefore, on the basis of previous knowledge and the new phytoplankton data collected, a photo guide of HABs species occurred in Apulia marine coastal areas (both toxin and high biomass producers) was developed with the following purposes:

- to support the technical staff of regional environmental agencies or students, as well as stakeholders and environmental managers in identifying, where possible, of phytoplankton species that may be harmful to human health or to productivity of the shellfish harvesting areas.
- to provide a practical approach which might facilitate the identification of phytoplankton organisms through morphological structures and characteristics visible mainly through optical light microscopy techniques (Bright field phase contrast, DIC and Epifluorescence)
- to provide an identification key of some genera that include HABs (Alexandrium, Dinophysis, Pseudo-nitzschia etc.) through a general schematic outlining of their major features.

Letteratura citata

Roselli L., Caroppo C., Bevilacqua S., Ciciriello P.C., Ungaro N., Vadrucci M.R. Harmful algae and pressure-impact relationship: noxious blooms and toxic microalgae occurrence from coastal waters of the Apulia region (Adriatic and Ionian Seas, Mediterranean). Submitted to Marine Environmental Research.

D.G.R n. 2014/2019. Intesa, ai sensi dell'articolo 8, comma 6, della legge 5 giugno 2003, n. 131 tra il Governo, le Regioni e le Province autonome di Trento e Bolzano concernente linee guida per l'applicazione del Regolamento (CE) 854/2004 e del Regolamento (CE) 853/2004 nel settore dei molluschi bivalvi.

AUTORI

Stefania D'Arpa (s.darpa@arpa.puglia.it; Autore di riferimento), Maria Rosaria Vadrucci (m.vadrucci@arpa.puglia.it), Pierangelo Cosimo Ciciriello, Elena Stanca, Anna Maria Pastorelli, Nicola Ungaro, Agenzia Regionale per la Prevenzione e Protezione Ambientale (ARPA Puglia), Corso Trieste 27, 70126.



Sessione Microalge III

Moderatore: Marina Montresor



Valutazione dell'effetto inibitorio di materiali legnosi modificati o trattati con impregnanti di nuova generazione sullo sviluppo della comunità microfitobentonica nella laguna di Grado (nord Adriatico)

V. Natali, T. Cibic

Qualsiasi substrato solido immerso in acqua marina o lagunare viene colonizzato dal biofilm microbico formato principalmente da batteri e diatomee. Per valutare come la formazione del biofilm risponda ad un possibile effetto inibitorio prodotto da materiali legnosi modificati o trattati con impregnanti, è stato condotto uno studio sperimentale nell'ambito del progetto DuraSoft finanziato dal programma interregionale Italia-Slovenia. A questo fine, sono stati preparati provini di abete bianco e abete rosso naturali per confrontarli con quelli opportunamente trattati con impregnanti di nuova formulazione (Silvanolin, rame etanoloammina), con prodotti già presenti sul mercato (CCB, rame-cromo-boro) o modificati termicamente per aumentarne la durevolezza. I provini di legno sono stati montati sui collettori e tenuti in acqua sempre alla stessa altezza (50 cm dal pelo dell'acqua, a prescindere dall'escursione di marea) mediante un sistema di boe di ancoraggio. In totale sono stati allestiti 24 collettori e posizionati in due siti della laguna di Grado in giugno 2021. I primi 12 sono stati posizionati nell'area "Schiusa" caratterizzata da elevato idrodinamismo, mentre i rimanenti 12 sono stati posizionati nell'area "Approdo" caratterizzata da condizioni idrodinamiche relativamente tranquille. In totale, sono stati immersi 144 provini, 72 per ciascuna area, 12 per ciascun tempo sperimentale, e recuperati dopo 6, 20 e 40 giorni per testare gli effetti a breve, medio e lungo periodo sull'abbondanza e composizione della comunità microalgale mediante diverse tecniche microscopiche (microscopia ottica, SEM, microscopia confocale a scansione laser). I risultati hanno evidenziato l'importanza dell'idrodinamismo dell'area: il minor idrodinamismo ha accelerato il processo di colonizzazione del biofilm microalgale, a prescindere dalla tipologia di trattamento applicato, con abbondanze cellulari fino a un ordine di grandezza maggiori. In condizioni di elevato idrodinamismo solo i trattamenti con Silvanolin hanno mantenuto l'efficacia inibente per tutta la durata dell'esperimento, invece l'efficacia degli altri trattamenti nell'inibire lo sviluppo del biofilm è scemata nel medio termine, dopo 20 giorni, o per effetto del dilavamento dei metalli pesanti oppure per il rilascio di composti nutritivi presenti negli impregnanti che hanno favorito la proliferazione microalgale. Focalizzandosi sull'effetto inibente dei diversi trattamenti, i generi *Nitzschia* e *Amphora* sono risultati più tolleranti, presentando abbondanze maggiori nei trattamenti rispetto al legno naturale. Altre sono risultate sensibili, scomparendo del tutto nei trattamenti con gli impregnanti, come il genere *Achnanthes*. In generale, gli impregnanti testati hanno indotto una serie di modificazioni alla comunità microalgale, rispetto al controllo sono state osservate diatomee di dimensioni inferiori, con frustuli malformati e strategie di vita che permettono di contrastare l'effetto tossico indotto dalle sostanze contenute negli impregnanti.

AUTORI

Vanessa Natali (vnatali@ogs.it), Tamara Cibic (tcibic@ogs.it; Autore di riferimento), Sezione di Oceanografia, Istituto Nazionale di Oceanografia e di Geofisica Sperimentale - OGS, via A. Piccard 54, 34151 Trieste, Italia.



Phytoplankton indicators for GES assessment: a case study in the N Adriatic Sea

F. Neri, M. Ubaldi, T. Romagnoli, S. Accoroni, C. Totti

Phytoplankton is considered as an optimal proxy to detect changes in the marine ecosystems, due to the fast turnover and the high sensitiveness to environmental conditions. For these reasons it is included as a descriptor for the assessment of the Good Environmental Status (GES) of pelagic habitats in the Marine Strategy Framework Directive, particularly at local scale. Effort must be done to test phytoplankton indicators at regional scale.

The aim of the study was to test several indicators using phytoplankton abundance data collected from 2008 to 2020 in 3 coastal stations of N Adriatic (Api, Senigallia and Portonovo).

Well-known indicators (i.e. Shannon diversity index and Pielou's evenness) were combined with newly ones (e.g. Rao index), to compare their efficiency. Multivariate statistical approaches (e.g. Non-Metric Multidimensional Scaling and clustering techniques) were used to look for patterns along the Mediterranean coasts. For each station, temporal changes were studied using the Local Contributions to Beta Diversity index (LCBD; a beta diversity measure) and the Importance Value Index (IVI) was calculated, revealing the taxa (genus) potentially responsible for the modification in the community structure.

This study will give information on the applicability of many indicators in the environmental assessment on a regional scale.

AUTORI

Francesca Neri (f.neri@pm.univpm.it; Autore di riferimento), Marika Ubaldi, Tiziana Romagnoli (t.romagnoli@univpm.it), Stefano Accoroni (s.accoroni@univpm.it), Cecilia Totti (c.totti@univpm.it), Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, via Brecce Bianche, 60131 Ancona, Italia.



Espressione genica durante la formazione di spore nella diatomea marina *Chaetoceros socialis*

A. Pelusi, L. Ambrosino, M. Miraldo, M.L. Chiusano, A. Rogato, M.I. Ferrante, M. Montresor

Molti organismi multicellulari ed unicellulari sono caratterizzati da periodi di quiescenza o dormienza durante i quali le attività metaboliche sono notevolmente ridotte. Tra le diatomee, microalghe unicellulari alla base delle reti trofiche acquatiche, diverse specie producono cellule dormienti (spore o 'resting cells') che possono sopravvivere nei sedimenti per lunghi periodi di tempo. Presenteremo i risultati di un primo studio sull'espressione genica durante il processo di formazione di spore in *Chaetoceros socialis*, una specie planctonica estremamente abbondante nei mari temperati. La formazione di spore è stata indotta da limitazione di azoto e l'esperimento di trascrittomico è stato condotto in condizioni di controllo (azoto non limitante) ed in tre momenti diversi durante la transizione da cellule vegetative a spore. Abbiamo rilevato *downregulation* di geni correlati alla fotosintesi e all'assimilazione dei nitrati, inclusi trasportatori di nitrati ad alta affinità (NTR). L'*upregulation* di altre vie cataboliche, come il ciclo dell'acido tricarbossilico, il ciclo del gliossilato e la beta-ossidazione, suggerisce che, durante il processo di formazione delle spore, questa diatomea potrebbe utilizzare i lipidi come fonte di energia. E' stata inoltre rilevata l'*upregulation* di una lipossigenasi e diverse aldeide deidrogenasi (ALDH) che supportano la presenza di ossilipine, metaboliti secondari con ruolo di mediatori chimici a livello intra-specifico. L'*upregulation* di geni coinvolti nella regolazione della dormienza in altri organismi (ad es. serina/treonina-protein chinasi TOR e il suo inibitore GATOR) rappresentano risultati interessanti che aprono la via a studi futuri. I risultati del nostro studio dimostrano che la transizione da una fase di crescita attiva ad una fase di dormienza è caratterizzata da marcati cambiamenti metabolici e forniscono evidenza della presenza di mediatori chimici che regolano la comunicazione fra le diverse cellule della popolazione.

AUTORI

Angela Pelusi (pelusi.angela@gmail.com; Autore di riferimento), Dipartimento di Ecologia Marina Integrata, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli; indirizzo presente: School of Earth and Environmental Sciences, Cardiff University, Main Building, Park Place, Cardiff, CF10 3AT, United Kingdom.

Luca Ambrosino (luca.ambrosino@szn.it), Marco Miraldo (marco.miralto@szn.it), Dipartimento Infrastrutture di Ricerca per le Risorse Biologiche Marine, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli, Italia.

Maria Luisa Chiusano (marialuisa.chiusano@unina.it), Dipartimento di Agraria, Università Federico II, Via Università 100, 80055 Portici e Dipartimento Infrastrutture di Ricerca per le Risorse Biologiche Marine, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli, Italia.

Alessandra Rogato (alessandra.rogato@ibbr.cnr.it), Istituto di Bioscienze e BioRisorse IBBR- CNR, Via Pietro Castellino 111, 80131 Napoli e Dipartimento di Ecologia Marina Integrata, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli, Italia.

Maria Immacolata Ferrante (mariella.ferrante@szn.it), Dipartimento di Ecologia Marina Integrata, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli e Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (OGS), Trieste, Italia.

Marina Montresor (marina.montresor@szn.it), Dipartimento di Ecologia Marina Integrata, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli, Italia.



Trait-based approach for rapid and effective characterization of phytoplankton patterns

L. Roselli, S. Bevilacqua, A. Terlizzi

Phytoplankton form the base of most aquatic food webs, accounting for approximately half of Earth's primary productivity. Also, such tiny microscopic organisms profoundly affect the biogeochemical cycling of many elements, such as carbon, nitrogen, and phosphorus. Phytoplankton determine the health of aquatic ecosystems, and have the potential, with harmful blooms, to reduce water quality and fishing yield. Because of the role that phytoplankton exert on the functioning of aquatic ecosystems, it is important to understand which factors govern the community assembly and dynamics. Human-induced pressures can alter phytoplankton community structure and, therefore, the need to understand the community reorganizations in the changing environment becomes even more pressing in the light of climate global change. For these reasons, phytoplankton assemblages are privileged descriptors of the ecological status of marine ecosystems and are regularly included in routine monitoring plans. Phytoplankton cell size and shape are key traits profoundly influence an organism's ecophysiological performance and evolutionary fitness, suggesting a link between morphology and diversity.

Trait-based approaches are increasingly used in phytoplankton ecology to explain and predict community organization along environmental gradients including human-induced ones. The high spatial and temporal variability of phytoplankton and the intrinsic difficulties of species identifications, however, combine in making reiterate assessments of this component of marine biota particularly demanding. Although with contrasting outcomes, coarse levels of taxonomic resolution (e.g., genus, family) or morpho-functional categories have been proposed to reduce identification efforts or to ease the analysis of phytoplankton assemblages for monitoring purposes. A major issue is that, in the absence of control for the loss of information associated to these alternative approaches, their application may lead to poor representations of genuine spatial and/or temporal patterns of assemblages in relation to natural and anthropogenic sources of variation. A new approach is proposed to reduce the efforts required to analyse phytoplankton assemblages that integrate morpho-functional classification of phytoplankton with the use of null models to estimate the consequent loss of information on species-level community patterns. Null models for information loss are built by randomly grouping the original species variables into a progressively decreasing number of groups, in order to identify the minimum number of aggregate variables needed to detect community patters as at species level. Aggregate variables are then defined as morpho-functional groups, by grouping species on the basis of a combination of morpho-functional traits, including general taxonomy, cell size, shape, elongation and complexity. The approach has been applied to several case studies investigating the response of phytoplankton assemblages in marine and transitional water ecosystems under different environmental settings in areas spanning the world's ocean, including coral atolls, mangroves, estuaries, coastal lagoons and inlets. The approach allowed obtaining parsimonious sets of morpho-functional groups, which were suitable to detect changes in phytoplankton assemblage structure as at species level in all case studies. Trait-based approaches to phytoplankton research and monitoring are crucial to shed light on processes underlying phytoplankton community assembly and dynamics in the face of global change. In this perspective, our framework incorporates cost-effectiveness, instances from traditional monitoring programs aiming at the detection of community patterns, and the current need for a deeper understanding of functional responses of phytoplankton to environmental drivers.

AUTORI

Leonilde Roselli (leonilde.roselli@szn.it, Autore di riferimento), Research Infrastructures for marine biological resources Department, Stazione Zoologica Anton Dohrn, Villa Comunale 1, 80121 Naples, Italy
 Stanislao Bevilacqua (sbevilacqua@units.it), Department of Life Sciences, University of Trieste, Via Licio Giorgieri 10, 34127 Trieste, Italy - CoNiSMA, Piazzale Flaminio 9, 00196 Roma, Italy



Antonio Terlizzi (antonio.terlizzi@szn.it), Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn, Villa Comunale 1, 80121 Naples, Italy - Department of Life Sciences, University of Trieste, Via Licio Giorgieri 10, 34127 Trieste, Italy - CoNiSMa, Piazzale Flaminio 9, 00196 Roma, Italy



Sessione Microalghe IV

Moderatore: Laura Pezzolesi



Dati preliminari su una microalga verde isolata dal Lido di Spina (Comacchio, Ferrara)

K. Sciuto, Sfriso A.A., M.A. Wolf, Sfriso A., E. Moschin, C. Munari, M. Mistri, I. Moro

Con il termine microalghe si intende un vasto gruppo di microrganismi fotossigenici appartenenti a taxa distinti, in grado di colonizzare svariati ambienti, anche quelli considerati estremi. Questi organismi possono, infatti, mettere in atto diversi adattamenti in risposta alle condizioni in cui si trovano e presentano, generalmente, un'elevata plasticità metabolica che gli permette di far fronte a cambiamenti anche repentini dell'ambiente circostante (Malcata *et al.*, 2018). Molte delle molecole prodotte dalle microalghe trovano impiego in vari ambiti di interesse umano, ad es. il settore agricolo, il cosmeceutico, il nutraceutico e della green energy (e.g., Levasseur *et al.*, 2020; Alvarez *et al.*, 2021; Kiran & Venkata Mohan, 2021), con il vantaggio che le sostanze prodotte dalle microalghe hanno generalmente nessuno o scarso impatto ambientale rispetto alle loro controparti di sintesi chimica. Inoltre, la produzione di microalghe richiede, solitamente, un limitato consumo di risorse e può essere condotta anche sfruttando sostanze di scarto (e.g. Shahid *et al.*, 2020).

In questo contesto è stato recentemente co-finanziato dal Fondo Sociale Europeo REACT EU – Programma Operativo Nazionale Ricerca e Innovazione 2014-2020, a titolarità del Ministero dell'Università e della Ricerca, un progetto dal titolo “Potenzialità biotecnologiche delle microalghe per la sostenibilità ambientale”, il cui scopo è analizzare diversi ceppi microalgali, caratterizzarli e valutarne la crescita in laboratorio, per poterli impiegare in diversi settori di interesse antropico. All'interno del progetto si considereranno sia ceppi microalgali già presenti presso collezioni pubbliche o private, che ceppi isolati ex-novo dall'ambiente, con particolare attenzione all'ambiente marino.

Sono stati, quindi, effettuati campionamenti nella zona del Lido di Spina, Comacchio (FE), andando a prelevare campioni d'acqua sia lungo la costa che da bacini d'acqua interni. Una volta in laboratorio, i campioni d'acqua sono stati processati e utilizzati per l'isolamento di diversi ceppi microalgali. Tra i ceppi isolati è risultata particolarmente interessante una microalga verde coccoide, temporaneamente denominata ceppo KS2, presente in un campione d'acqua prelevato da uno dei bacini interni, della cui caratterizzazione verranno qui presentati i primi dati. La microalga è stata sottoposta ad un approccio di tipo polifasico, comprendente analisi molecolari basate sul gene 18S rRNA e sullo spaziatore intergenico trascritto 18S-28S ITS rRNA, analisi morfologiche ed ultrastrutturali, mediante microscopia ottica ed elettronica (SEM e TEM), e analisi biochimiche (es. pigmenti, polisaccaridi, proteine), nonché la valutazione della crescita del microorganismo in condizioni di laboratorio.

Letteratura citata

- Alvarez A.L., Weyers S.L., Goemann H.M., Peyton B.M., Gardner R.D. (2021). Microalgae, soil and plants: A critical review of microalgae as renewable resources for agriculture. *Algal Research*, 54: Article 102200.
- Kiran B.R., Venkata Mohan S. (2021). Microalgal Cell Biofactory - Therapeutic, Nutraceutical and Functional Food Applications. *Plants*, 10: Article 836.
- Levasseur W., Perré P., Pozzobon V. (2020). A review of high value-added molecules production by microalgae in light of the classification. *Biotechnology Advances* 41: Article 107545.
- Malcata F.X., Sousa Pinto I., Guedes A.C. [Eds] (2018). Marine macro-and microalgae: an overview. CRC Press, Boca Raton, Florida, USA. 342 pp.
- Shahid A., Malik S., Zhu H., Xu J., Nawaz M.Z., Nawaz S., Asraful Alam M., Mehmood M.A. (2020). Cultivating microalgae in wastewater for biomass production, pollutant removal, and atmospheric carbon mitigation; a review. *Science of the Total Environment*, 704: Article 135303.

AUTORI

Katia Sciuto (katia.sciuto@unife.it; Autore di riferimento), Università di Ferrara, Dipartimento di Scienze Chimiche, Farmaceutiche e Agrarie, via L. Borsari 46, 44121 Ferrara.



Andrea A. Sfriso, Università di Ferrara, Dipartimento di Scienze Chimiche, Farmaceutiche e Agrarie, via L. Borsari 46, 44121 Ferrara.

Marion A. Wolf, Università di Ferrara, Dipartimento di Scienze Chimiche, Farmaceutiche e Agrarie, via L. Borsari 46, 44121 Ferrara.

Adriano Sfriso, Università Ca' Foscari di Venezia, Dipartimento di Scienze Ambientali, Informatica e Statistica, via Torino 155, 30172 Venezia-Mestre.

Emanuela Moschin, Università di Padova, Dipartimento di Biologia, via U. Bassi 58/B, 35131 Padova.

Cristina Munari, Università di Ferrara, Dipartimento di Scienze Chimiche, Farmaceutiche e Agrarie, via L. Borsari 46, 44121 Ferrara.

Michele Mistri, Università di Ferrara, Dipartimento di Scienze Chimiche, Farmaceutiche e Agrarie, via L. Borsari 46, 44121 Ferrara.

Isabella Moro, Università di Padova, Dipartimento di Biologia, via U. Bassi 58/B, 35131 Padova.



Effect of salinity on growth and microcystin production by the cyanobacterium *Planktothrix rubescens*

M. Ubaldi, S. Bacchiocchi, M. Siracusa, F. Neri, T. Romagnoli, C. Totti, S. Accoroni

The cyanobacterium *Planktothrix rubescens* is a common producer of cyanotoxins (mainly microcystins), which forms blooms in freshwaters. During algal blooms, microcystins can reach the sea and can be accumulated in bivalve molluscs with risks for consumption. In northern coast of Puglia (near the Lesina and Varano lakes), were found mussels that accumulated microcystins produced by *P. rubescens* at concentrations higher than tolerable daily intake (De Pace et al., 2014). In this study we tested the tolerance and microcystin production of *P. rubescens* to increasing salinity. One strain of *P. rubescens* was isolated from the lake of Castreccioni (Cingoli, MC) and exposed for one month to different salinities (0, 10, 20, 30) in batch cultures. Cell abundances and growth rate were obtained by optical density (OD650 e OD750) at the spectrophotometer and by counts through light microscope. Both methods showed that *P. rubescens* is unable to survive for more than 96 hours at salinity equal to or greater than 10. Anyway, the inhibition is not immediate, as for the first 4 days the abundances remained similar to the control. This implies that once the cells have reached the coasts, they remain vital enough to allow filter-feeding organisms to accumulate cells and toxins in their tissues. At the IZSUM, toxins were quantified in the cultures both in intracellular compartment and in culture medium (dissolved form) using LC-MS/MS. The most found microcystin was MC-RR desmethylated. Higher intracellular concentrations of the microcystins were observed at lower salinities (0 and 10), while the amount of the dissolved form was higher at salinities 20 and 30. Our results suggest that salinity stress could cause plasmolysis and release of toxins as already found by Rosen et al. (2018) for cyanobacteria. Monitoring of cyanotoxins in sea water and in bivalve molluscs is important for the public health.

Letteratura citata

- De Pace R., Vita V., Bucci M., Gallo P., Bruno M. (2014). "Microcystin contamination in sea mussel farms from the italian southern adriatic coast following cyanobacterial blooms in an artificial reservoir". In: Journal of Ecosystems 2014.
 Barry H., Keith A., Jennifer L., Katherine N., James M., Brett D., Sarena S. (2018). "Understanding the effect of salinity tolerance on cyanobacteria associated with a harmful algal bloom in Lake Okeechobee, Florida". 2018-5092. US Geological Survey.

AUTORI

Marika Ubaldi (m.ubaldi@staff.univpm.it), Autore di riferimento), Francesca Neri (f.neri@pm.univpm.it), Tiziana Romagnoli (tromagnoli@staff.univpm.it), Stefano Accoroni (s.accoroni@staff.univpm.it), Cecilia Totti (c.totti@univpm.it), Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, via Brecce Bianche, 60131 Ancona, Italy.
 Simone Bacchiocchi (s.bacchiocchi@izsum.it), Melania Siracusa (m.siracusa@izsum.it), Istituto Zooprofilattico Sperimentale Umbria e Marche "Togo Rosati", Via G. Salvemini, 1, 06126 Perugia, Italy.



Moschin Potential of a green algal strain collected from the Sarno River mouth for heavy metal bioremediation

L. Barra, S. Balzano, M. Moros, A. Smerilli, E. Cavalletti, I. Percopo, P. Chiaiese, A. Sardo

Heavy Metals (HMs) can induce both chronic and acute harmful effects on marine and freshwater biota and their presence in soils, sediments, and waters has been well documented (Yabanli et al. 2015). In this context, eukaryotic microalgae are actually considered as suitable microorganisms for bioremediation, since they (1) do not require organic substrates neither clean water to achieve fast growth, (2) they are able to uptake HMs and other pollutants from different environments, including wastewaters and contaminated sediments, and (3) they compartmentalize pollutants in vacuoles or other organelles minimizing their detrimental effects (Balzano et al., 2020). The Sarno River is a 30 km long freshwater stream located in Southern Italy that flows into the Gulf of Naples. Partially untreated industrial wastewaters, deriving from the tanning and canning industries, have been discharged directly to Sarno River or its tributaries for several decades. This led to significant pollution in hydrocarbons and HMs along the river as well as in the seawater basin in proximity of its mouth (Montuori et al., 2013). We aimed at determining the ability of a green microalgal strain, isolated from Sarno River Mouth, *Tetraselmis convolutae* strain BS14, to grow in artificially polluted seawater and to accumulate selected HMs within its biomass. We performed the molecular characterization of the species, the inhibition growth rate, and the analysis of Ion Coupled Plasma-Mass Spectrometry (ICP-MS) in presence of Cadmium (Cd), Lead (Pb) and Zinc (Zn). The metals were analyzed one at time, in order to investigate the potentiality of this green algal strain in the phyco-remediation process for future biotechnological application. *T. convolutae* BS14 performed a reasonably fast growth in the presence of cadmium (up to 50 µM), zinc (20 µM), and lead (50 µM) for seven days. Cell surface area slightly increased in the presence of HMs and cells were found to incorporate part of the HMs. The present study indicates the potential of *T. convolutae* BS14 for the bioremediation of HM contaminated waters.

Letteratura citata

- Balzano S., Sardo A., Blasio M., Chahine T. B., Dell'Anno F., C. Sansone C. and Brunet C. (2020). Microalgal Metallothioneins and Phytochelatins and Their Potential Use in Bioremediation. *Frontiers in Microbiology*, 11.
- Montuori, P., P. Lama, S. Aurino, D. Naviglio and M. Triassi (2013). Metals loads into the Mediterranean Sea: estimate of Sarno River inputs and ecological risk. *Ecotoxicology* 22(2): 295-307.
- Yabanli, M., Y. Coskun and B. Oz (2015). Detection of heavy metal concentrations in three freshwater fish species in lake marmara (western anatolia) and heavy metal human health risk assessment."Fresenius Environmental Bulletin 24(12): 4273-4278.

AUTORI

- Lucia Barra (lucia.barra@szn.it: Autore di riferimento), Dipartimento di Biotecnologie Marine Ecosostenibili (BLUBIO) Stazione Zoologica Anton Dohrn, Calabria Marine Centre Contrada Torre Spaccata 87071 Amendolara (Cs), Italia.
- Sergio Balzano (sergio.balzano@szn.it), Dipartimento di Biotecnologie Marine Ecosostenibili (BLUBIO) Stazione Zoologica Anton Dohr Napoli (SZN), Giardini del Molosiglio via Acton 55, 80133 Napoli, Italia.
- Maria Moros Caballero (mamoros@unizar.es), Instituto de Nanociencia y Materiales de Aragón (INMA) Esquillor 15, 50018 Zaragoza, Spagna.
- Pasquale Chiaiese (pasquale.chiaiese@unina.it), Dipartimento di Agraria, Università degli Studi di Napoli Federico II, *Parco Gussone*, via Università 133 Portici, Napoli, Italia.
- Isabella Percopo (percopo@szn.it), Dipartimento di infrastrutture per la ricerca e le risorse in biologia marina (RIMAR), Stazione Zoologica Anton Dohrn, Villa Comunale 80121 Napoli, Italia.
- Arianna Smerilli (arianna.smerilli@szn.it); Elena Cavalletti (elena.cavalletti@szn.it), Angela Sardo (angela.sardo@szn.it), Dipartimento di Biotecnologie Marine Ecosostenibili (BLUBIO) Stazione Zoologica Anton Dohrn, Giardini del Molosiglio via Acton 55, 80133 Napoli, Italia.



Investigation of potentially toxic metabolites in a newly isolated freshwater cyanobacterium of the family Pseudanabaenaceae through a genomic approach

M. Simonazzi, L. Pezzolesi, T.K. Shishido, D.P. Fewer, R. Pistocchi

In the past decades, the proliferation of cyanobacteria in freshwater ecosystems has intensified, possibly as a consequence of climate changes and eutrophication. The ever-increasing presence of these phototrophic organisms in water bodies for human use (e.g. drinking water, recreational activities) is considered as an emerging health threat, since some cyanobacteria are able to produce harmful toxins, named after them cyanotoxins. So far, the discovery of cyanotoxins has been investigated after a human or animal intoxication event occurred, through a combination of chemical and biological analyses carried out on bloom samples or cultured organisms isolated from their natural environment. In recent years, the advances in genome sequencing and analysis have broaden our understanding and knowledge of prokaryotes such as cyanobacteria by in-depth investigations at genome level, thus expanding the potential for the discovery of unknown metabolites, including new toxic compounds. In the present work, a cyanobacterium belonging to the family of Pseudanabaenaceae was isolated from a small lake that provides water for the local drinking water plant (Rimini, Emilia-Romagna, Italy). Aqueous cyanobacterial extracts were tested by *Artemia* sp. bioassay, showing high mortality rates of the individuals within 24h (98-100%). Chemical analyses (LC-MS) on methanolic extracts, and PCR amplification of genes involved in the synthesis of various cyanotoxins did not give any results, thus the production of known cyanotoxins in the strain was excluded. The genome of the strain was then sequenced by Illumina MiSeq, obtaining a completeness of 99.64%. Genome analyses confirmed the absence of known cyanotoxins gene clusters, while revealed the presence of a region responsible for the synthesis of polyketides (i.e. trans-AT PKS), a highly diverse class of compounds both in terms of structure (i.e. alkaloid, macrolides, polyethers, aromatics) and biological activities (i.e. antibiotic, anticancer, antifungal), some of which considered involved in toxicity. The end product of this region has not been detected yet, and further analyses are needed to elucidate its structure and potential toxicological characteristics. Since the toxic effects were observed in aqueous extracts of the cyanobacterium, potential implication for the safety of drinking waters should be taken into consideration.

AUTORI

Mara Simonazzi, (mara.simonazzi2@unibo.it; Autore di riferimento), Department of Biological, Geological and Environmental Sciences-BiGeA, University of Bologna, Via Sant'Alberto, 163, 48123 Ravenna, Italia.

Laura Pezzolesi, Rossella Pistocchi, Department of Biological, Geological and Environmental Sciences-BiGeA, University of Bologna, Via Sant'Alberto, 163, 48123 Ravenna, Italia.

Tânia K. Shishido, David P. Fewer, Department of Microbiology, University of Helsinki, Viikinkaari 5, 00790 Helsinki, Finlandia.



Organelle Phylogenomics unveil the diverging evolution of 6 Lineages in *Galdieria sulphuraria* (Cyanidiophyceae).

M. Iovinella, C. Ciniglia, S.J. Davis

Exploration of life in extreme environments allows the discovery of intriguing organisms with extraordinary biotechnological potential. An example of extreme environments is represented by hot springs, where harsh conditions ($\text{pH} < 1$; Temperature $> 50^\circ\text{C}$; high concentrations of metals) are prohibitive for most living organisms, except for archaea, bacteria and a few eukaryotes like the unicellular red alga *Galdieria* (Iovinella et al., 2018). Phylogenetic analysis based on a few plastids and nuclear genes highlighted the intricate genetic structure of *Galdieria* and the hypothesis of diverging clades within the *G. sulphuraria* species (Toplin et al., 2008, Hsieh et al., 2015). To resolve enigmatic relationships between lineages, we used plastid and mitochondrial genome-scale data obtained from numerous strains from around the world. The resulting phylogenomic analysis identified: i) the divergence of each of the mitochondrial and plastid genomes into the same six clear lineages; ii) the independent evolution of the lineages; iii) the incongruent interlineages relationships between the genomes. Differential evolutionary pressure between the strains and the genomes were also highlighted by synonymous and non-synonymous substitutions.

Letteratura citata

Iovinella M, Eren A, Pinto G, Pollio A, Davis SJ, Cennamo P, et al. Cryptic dispersal of Cyanidiophytina (Rhodophyta) in non-acidic environments from Turkey. *Extremophiles* [Internet]. 2018 Sep;22(5):713–23. Available from: <http://dx.doi.org/10.1007/s00792-018-1031-x>

Toplin JA, Norris TB, Lehr CR, McDermott TR, Castenholz RW. Biogeographic and phylogenetic diversity of thermoacidophilic cyanidiales in Yellowstone National Park, Japan, and New Zealand. *Appl Environ Microbiol* [Internet]. 2008 May;74(9):2822–33. Available from: <http://dx.doi.org/10.1128/AEM.02741-07>

Hsieh CJ, Zhan SH, Lin Y, Tang SL, Liu SL. Analysis of rbcL sequences reveals the global biodiversity, community structure, and biogeographical pattern of thermoacidophilic red algae (Cyanidiales). *J Phycol* [Internet]. 2015 Aug;51(4):682–94. Available from: <http://dx.doi.org/10.1111/jpy.12310>

AUTORI

Manuela Iovinella (manuela.iovinella@unicampania.it; Autore di riferimento), Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Caserta “L.Vanvitelli”, Via Vivaldi 43, 81100 Caserta, Italia.

Claudia Ciniglia (claudia.ciniglia@unicampania.it), Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Caserta “L.Vanvitelli”, Via Vivaldi 43, 81100 Caserta, Italia.

Seth J. Davis (seth.davis@york.ac.uk), Department of Biology, University of York, Wentworth Way, York YO10 5DD



Sessione Microalge V

Moderatore: Chiara Pennesi



***Halomicronema metazoicum*, la storia di un cianobatterio, del suo ambiente e dei suoi vicini**

M. Mutualipassi, V. Mazzella, N. Ruocco, V. Zupo

I cianobatteri svolgono ruoli ecologici chiave in numerosi ambienti e mostrano un ampio adattamento ad innumerevoli differenti habitat. Molte specie producono molecole bioattive via secrezione costitutiva, in altri casi, alcune molecole prodotte sono invece metaboliti secondari attivati dal grazing (solitamente Wound-Activated Compounds). Sia le molecole costitutivamente espresse che quelle Wound-activated possono avere anche funzione di messaggio informazionale di tipo chimico (infochimico). I nostri studi si sono focalizzati su un particolare cianobatterio del gruppo *Phormidium*-like, *Halomicronema metazoicum* C.Caroppo, P.Pagliara & P.Albertano, 2012, che possiede caratteristiche interessanti da un punto di vista ecologico e biotecnologico. E' stato identificato per la prima volta in simbiosi con la spugna *Petrosia ficiformis* (Poiret, 1789), con probabili ruoli antigrazing e azotofissazione (Pagliara et al., 2020). Nel 2016 è stato da noi isolato come colonia non associata a spugne e ritrovato come epifita di foglie di *Posidonia oceanica* (L.) Delile, 1813 (Ruocco et al., 2018). *Halomicronema metazoicum* è stato inoltre isolato nella colonna d'acqua testimoniando il comportamento quantomeno meroplanctonico della specie, suggerendo inoltre ruoli ecologici complessi in questi ambienti. Test comportamentali su alcuni invertebrati di fauna vagile hanno evidenziato come le specie campionate dalla prateria evitassero le aree arricchite di bouquet odorosi prodotti da *H. metazoicum*, riconoscendo gli infochimici del cianobatterio e la sua aggressività chimica (Zupo et al., 2016). Alcune evidenze suggeriscono come i bloom bentonici e planctonici possano essere dettati da fenomeni di allelopatia dettati da infochimici e quindi da comunicazione tra specie (Mutalipassi et al., 2021). Campionamenti hanno evidenziato come la presenza di alcuni cianobatteri meroplanctonici nella colonna d'acqua o nel benthos siano inversamente correlati con presenza o assenza di altri organismi, come *Ostreopsis ovata* Y.Fukuyo 1981, negli stessi comparti. Test da noi effettuati, a sostegno di questa tesi, dimostrano come molecole bioattive prodotte in certe condizioni dal cianobatterio provochino l'incistamento immediato di *O. ovata* senza però causarne la morte. Tali molecole hanno mostrato elevata tossicità su plutei di *Arbacia lixula* (Linnaeus, 1758) e *Paracentrotus lividus* (Lamarck, 1816), nonché su vari stadi di *Brachionus plicatilis* (Müller, 1786) (Mutalipassi et al., 2019, 2022). I mezzi di coltura sono stati anche testati su copepodi parassiti nonché su alcune linee cellulari (tumorali e non) umane, al fine di valutare il potenziale biotecnologico delle molecole costitutivamente prodotte. In conclusione, gli studi condotti dimostrano da un lato l'enorme ruolo svolto dai cianobatteri nelle comunità litorali, con un'ampia diversità di processi biochimici attivi e una vasta gamma di interazioni con altre specie; dall'altro lato dimostrano come, nonostante la lunga tradizione di screening e separazione di molecola bioattive, la biodiversità chimica e di specie dei cianobatteri è tuttora sottostimata, suggerendo grandi prospettive future per la drug discovery.

Letteratura citata

- Mutalipassi, M., Mazzella, V., Glaviano, F., and Zupo, V. (2022). Constitutive and activated toxigenic activity of *Halomicronema metazoicum* (Cyanoprokaryota, Cyanophyta). Mar. Ecol. e12697. doi: 10.1111/maec.12697.
- Mutalipassi, M., Mazzella, V., Romano, G., Ruocco, N., Costantini, M., Glaviano, F., et al. (2019). Growth and toxicity of *Halomicronema metazoicum* (Cyanoprokaryota, Cyanophyta) at different conditions of light, salinity and temperature. Biol. Open 8. doi: 10.1242/bio.043604.
- Mutalipassi, M., Riccio, G., Mazzella, V., Galasso, C., Somma, E., Chiarore, A., et al. (2021). Symbioses of Cyanobacteria in marine environments: Ecological insights and biotechnological perspectives. Mar. Drugs 19. doi: 10.3390/md19040227.
- Pagliara, P., Barca, A., Verri, T., and Caroppo, C. (2020). The marine sponge *Petrosia ficiformis* harbors different cyanobacteria strains with potential biotechnological application. J. Mar. Sci. Eng. 8. doi: 10.3390/JMSE8090638.
- Ruocco, N., Mutualipassi, M., Pollio, A., Costantini, S., Costantini, M., and Zupo, V. (2018). First evidence of *Halomicronema metazoicum* (Cyanobacteria) free-living on *Posidonia oceanica* leaves. PLoS One 13, e0204954. doi: 10.1371/journal.pone.0204954.



Zupo, V., Mutualipassi, M., Fink, P., and Di Natale, M. (2016). Effect of Ocean Acidification on the communications among invertebrates mediated by plant-produced Volatile Organic Compounds. *Glob. Ecol. Biogeogr.* 1, 012–018. doi: 10.17352/gie.000002.

AUTORI

Mirko Mutualipassi (mirko.mutalipassi@szn.it; Autore di riferimento), Stazione Zoologica Anton Dohrn, Calabria Marine Centre, Contrada Torre Spaccata, Amendolara, 87071 Cosenza, Italia.
Valerio Mazzella, (valerio.mazzella@szn.it), Stazione Zoologica Anton Dohrn, Villa Comunale, 80121, Napoli, Italia.
Nadia Ruocco, (nadia.ruocco@szn.it), Stazione Zoologica Anton Dohrn, Calabria Marine Centre, Contrada Torre Spaccata, Amendolara, 87071 Cosenza, Italia.
Valerio Zupo, (valerio.zupo@szn.it), Stazione Zoologica Anton Dohrn, Ischia Marine Centre, Via Porto, 127, Ischia 80077 Napoli, Italia.



Induzione di stadi di resistenza nella diatomea marina *Thalassiosira rotula*

I. Orefice, V. Di Dato, G. Romano

Molte specie fitoplanctoniche, ed in particolare le diatomee, adottano la strategia di formare stadi di resistenza in risposta a condizioni sfavorevoli in termini di nutrienti, luce e temperatura, consentendo la sopravvivenza delle cellule e contribuendo al loro successo ecologico (McQuoid e Hobson, 1996). Si tratta infatti di uno stato reversibile dal quale le cellule sono in grado di riprendere la crescita vegetativa quando i fattori ambientali diventano nuovamente favorevoli. I meccanismi che sottendono la formazione di stadi di resistenza possono differire all'interno di uno stesso gruppo di organismi e anche nell'ambito dello stesso genere o specie; quindi, è necessaria una precisa descrizione di tali meccanismi a livello di specie. Abbiamo studiato l'induzione di forme di resistenza nella diatomea centrica *Thalassiosira rotula*, valutando le condizioni che ne inducono la formazione e la germinazione, con lo scopo finale di indentificare i metaboliti responsabili con possibile interesse per scopi biotecnologici. *T. rotula* è una specie cosmopolita costiera, presente nel Golfo di Napoli, che spesso domina la comunità del fitoplancton per le sue grandi dimensioni e l'elevata concentrazione. Abbiamo stabilito le condizioni per indurre le cellule di *T. rotula* ad entrare in uno stato di dormienza valutando diverse concentrazioni di nutrienti (azoto, silice e ferro), di intensità luminosa, densità cellulare e livelli di ossigenazione. Una bassa disponibilità di azoto, una bassa densità cellulare iniziale e un buon livello di ossigenazione sono risultati i principali fattori che determinano l'induzione dello stato di dormienza in questa specie di diatomee. La germinazione è stata testata dopo diversi tempi di conservazione, dimostrando che queste cellule dormienti hanno una capacità di resistenza almeno fino a due mesi. L'osservazione al microscopio ottico e a fluorescenza ha mostrato che le forme di resistenza di *T. rotula* sono morfologicamente simili a quelle vegetative ma con un frustulo più spesso, un citoplasma condensato e addossato alle pareti e un livello di pigmentazione inferiore.

Questo studio preliminare fornisce la base per successivi studi volti ad identificare i meccanismi molecolari coinvolti nella formazione di stadi di resistenza in questa specie. A questo scopo, intendiamo valutare l'espressione differenziale dei geni coinvolti tramite analisi trascrittometrica comparativa. Verrà inoltre studiato il possibile coinvolgimento di modificazioni epigenetiche nella transizione tra i due stadi.




Figura 1 Immagini della diatomea *Thalassiosira rotula* in stato vegetativo (A) and in stadio di resistenza (B) ottenute al microscopio ottico.

Letteratura citata

- Di Costanzo F., Di Dato V., van Zyl L.J., Cutignano A., Esposito F., Trindade M., Romano G. (2021). Three novel bacteria associated with two centric diatom species from the Mediterranean Sea, *Thalassiosira rotula* and *Skeletonema marinoi*. International Journal of Molecular Science, 22(24): 13199.
McQuoid, M.R. and Hobson, L.A. (1996). Diatom resting stages. Journal of Phycology, 32: 889-902.

AUTORI



RIUNIONE SCIENTIFICA ANNUALE
Amendolara, 11-12 novembre 2022



Ida Orefice (idaorefice@szn.it), Valeria Di Dato (valeria.didato@szn.it; Autore di riferimento), Giovanna Romano (giovanna.romano@szn.it), Sustainable Marine Biotechnology Department, Stazione Zoologica Anton Dohrn, via Ammiraglio Ferdinando Acton 55, 80133 Napoli, Italia.



Production of the biopolymer polyhydroxybutyrate (PHB) from the Chlorophyceae *Desmodesmus communis*

G. Zoffoli, C. Samorì, M. Simonazzi, R. Pistocchi, L. Pezzolesi

The sustainability of plastic materials is a major focus of basic research, industry, government, and society to find substitutes to petroleum-based and non-biodegradable conventional plastics. Polyhydroxyalkanoates (PHAs) are promising alternatives that accumulate as energy and carbon storage material in various microorganisms, including bacteria and microalgae, being biodegradable and suitable for a wide variety of applications. The most prevalent and well-characterized biopolymer is the polyhydroxybutyrate (PHB), which belongs to the short-chain PHAs.

The present study was designed to evaluate algae-based PHB production using the Chlorophyceae *Desmodesmus communis* in two-phase nutritional mode of cultivation, namely a phototrophic growth phase (PGP) and a mixotrophic stress phase (MSP). Microalgal PHB accumulation was in fact driven by phosphate constraint, and organic carbon (i.e. sodium acetate) supply.

The highest PHB productivity (0.11 g PHB/g biomass/d; 0.015 g PHB/L/d), corresponding to 32.1% w/w of intracellular PHB, was observed for *D. communis* after 3 days of cultivation under mixotrophic conditions (e.g., low light, phosphorus-free medium, 1 g/L of sodium acetate). A scaled-up cultivation (10 L) was set up to evaluate the PHB yield in a semi-continuous system in relation to the biomass main composition. After 10 days of cultivation, a content of 37.9% w/w of proteins, 24.1% w/w of polysaccharides, 23.1% w/w of PHB, and 9.0% w/w of lipids was reached. As bacteria are traditionally the principal PHB producers, preliminary tests performed with axenic cultures were performed to understand the contribution of *D. communis* associated microbiota to the PHB production. Results highlighted as the achievement of a biomass dry weight of about 0.2-0.3 g/L is necessary to optimize PHB production from *D. communis* cells, resulting in a productivity up to 2-times higher than those reported for cyanobacterial species (Simonazzi et al., 2021) and comparable to the maximum value obtained for other Scenedesmaceae (García et al., 2020; Kumari et al., 2022). In addition, considering *D. communis* ability to bioremediate wastewaters, its exploitation in a circular economy perspective could be promising. Further investigations should be addressed to better understand the physiological mechanisms that allow this species to produce PHB, considering the importance of the associated microbiota.

Letteratura citata

- García, G., Sosa-Hernández, J.E., Rodas-Zuluaga, L.I., Castillo-Zacarías, C., Iqbal, H., Parra-Saldívar, R., 2020. Accumulation of PHA in the Microalgae *Scenedesmus* sp. under Nutrient-Deficient Conditions. Polymers 13, 131.
- Kumari, P., Kiran, B.R., Mohan, S.V., 2022. Polyhydroxybutyrate production by Chlorella sorokiniana SVMIICT8 under Nutrient-deprived mixotrophy. Bioresource Technology 354, 127135. <https://doi.org/10.1016/j.biortech.2022.127135>
- Simonazzi, M., Pezzolesi, L., Galletti, P., Gualandi, C., Pistocchi, R., De Marco, N., Paganelli, Z., Samorì, C., 2021. Production of polyhydroxybutyrate by the cyanobacterium cf. *Anabaena* sp. Int. J. Biol. Macromol. 191, 92–99.

AUTORI

Giorgia Zoffoli, Dipartimento di Scienze Biologiche, Geologiche ed Ambientali (BiGeA), University of Bologna, via Sant'Alberto 163, 48123, Ravenna, Italia.

Chiara Samorì, Centro Interdipartimentale di Ricerca Industriale Fonti Rinnovabili, Ambiente, Mare ed Energia (CIRI-FRAME), University of Bologna, via Sant'Alberto 163, 48123, Ravenna, Italia; Dipartimento di Chimica "Giacomo Ciamician", University of Bologna, via Sant'Alberto 163, Ravenna, Italia.

Mara Simonazzi (mara.simonazzi2@unibo.it), Dipartimento di Scienze Biologiche, Geologiche ed Ambientali (BiGeA), University of Bologna, via Sant'Alberto 163, 48123, Ravenna, Italia.



RIUNIONE SCIENTIFICA ANNUALE
Amendolara, 11-12 novembre 2022



Rossella Pistocchi, Laura Pezzolesi (laura.pezzolesi@unibo.it; Autore di riferimento), Dipartimento di Scienze Biologiche, Geologiche ed Ambientali (BiGeA), University of Bologna, via Sant'Alberto 163, 48123, Ravenna, Italia; Centro Interdipartimentale di Ricerca Industriale Fonti Rinnovabili, Ambiente, Mare ed Energia (CIRI-FRAME), University of Bologna, via Sant'Alberto 163, 48123, Ravenna, Italia.



Indice degli autori

Accoroni S.	27, 34	Di Dato V.	41
Ambrosino L.	28	Drigo F.	18
Auciello C.	2, 20	Duarte Lopes A.	3
Bacchicocchi S.	34	Dupont G.	22
Balzano S.	22, 35	Falace A.	13
Bandelj V.	13	Fanelli G.	7
Barra L.	35	Ferrante M.I.	28
Bartual A.	9	Fewer D.P.	36
Bentivogli R.	4	Fondo J.	4
Bevilacqua S.	29	Gianni F.	13
Blasio M.	22, 35	Iovinella M.	2, 20, 37
Bolpagni R.	16, 17	Kaleb S.	13
Bottalico A.	11	Lenzo D.	9
Caputo N.	4	Marieschi M.	16, 17
Carbone M.	10	Marotta P.	22
Cavalletti E.	35	Mastrototaro F.	11
Cecere E.	7	Mazzella V.	39
Chiaiese P.	35	Mazzuca S.	10
Chimienti G.	11	Miraldo M.	28
Chiusano M.L.	28	Mistri M.	32
Cibic T.	26	Molinari G.	16, 17
Ciciriello P.C.	24	Mollo E.	10
Ciniglia C.	2, 20, 37	Mollo L.	18
Colangelo M.A.	9	Montresor M.	28
Costantini F.	4	Moro I.	3, 32
Cuevas B.	9	Moros M.	35
Custódio L.	3	Moschin E.	32
Cutignano A.	22	Munari C.	32
D'Arpa S.	24	Mutalipassi M.	39
Davis S.J.	37	Natali V.	26
de Klerk G.V.	22	Neri F.	27, 34
DeLuca P.	22	Nisticò D.M.	10
Denti G.	7	Norici A.	18
Descourvières E.	13	Oliva D.	10
De Stefano M.	2, 20	Orefice I.	41



Orlando-Bonaca M.	13	Schiaparelli S.	3
Ossò V.	10	Sciuto K.	32
Papa L.	7	Sfriso A.	32
Papirio S.	22	Sfriso A.A.	32
Pasteris A.	9	Shishido T.K.	36
Pastorelli A.	24	Simonazzi M.	36, 43
Pelusi A.	28	Siracusa M.	34
Pennesi C.	2, 20	Smerilli A.	35
Percopo I.	35	Stanca E.	24
Petrocelli A.	7	Terlizzi A.	29
Pezzolesi L.	4, 9, 36, 43	Torelli A.	17
Piro A.	10	Totti C.	27, 34
Pistocchi R.	4, 9, 36, 43	Trancone G.	22
Rabaoui L.	20	Trentin R.	3
Rogato A.	28	Tursi A.	11
Romagnoli T.	27, 34	Ubaldi M.	27, 34
Romano G.	41	Úbeda B.	9
Roselli L.	29	Ungaro N.	24
Rubino F.	7	Vadrucci M.R.	24
Ruocco N.	39	Wolf M.A.	32
Samorì C.	43	Zoffoli G.	43
Sardo A.	22, 35	Zubía E.	9
Savi S.	16	Zupo V.	39
Scarcelli F.	10		